

VOL. XII,
OLD SERIES.

PHILADELPHIA, SEPTEMBER, 1898.

VOL. I. No. 9.
NEW SERIES.

PUBLISHED IN CONJUNCTION WITH
Sajous's Annual and Analytical Cyclopædia of Practical Medicine.


PRICE OF THIS JOURNAL, SEPARATE, \$5.00 PER YEAR.

THE
Monthly Cyclopædia
OF
Practical Medicine
AND
UNIVERSAL MEDICAL JOURNAL.

EDITED BY

CHARLES E. de M. SAJOUS, M.D.,

PHILADELPHIA.



LEADING ARTICLES: "Amenorrhœa." "Gout." "Sterility." "Yellow Fever."

THE F. A. DAVIS CO., Publishers.

Office: 1914 and 1916 Cherry Street, Philadelphia, Pa., U. S. A.

BRANCH OFFICES: New York, U. S. A., 117 W. Forty-Second St.; Chicago, Ill., U. S. A., 9 Lakeside Building.

Exchanges and editorial matter to be sent to 2043 Walnut Street, Philadelphia, Pa.
All letters concerning subscriptions, removals, advertisements, etc., to be sent to the Publishers.
This Journal should be carefully preserved for binding as an additional volume of the Cyclopædia
at the end of each year.

"The general resemblance of milk modified by the PEPTOGENIC MILK POWDER to the infant's natural supply, is a very striking one." R. H. CHITTENDEN.

"When properly prepared, the resultant, so-called 'humanised milk', (prepared with PEPTOGENIC POWDER), presents the albuminoids in a minutely coagulable and digestible form; has an alkaline reaction; contains the proper proportion of salts, milk sugar, and fat; is not bitter in taste, being but partially peptonised, and in appearance as well as chemical composition resembles human milk." LOUIS STARR.

PACKER'S TAR SOAP.

VEGETABLE OILS, PINE-TAR,
AND GLYCERINE.



PURE. ANTISEPTIC.
EMOLLIENT.

THE STANDARD.—Used and recommended by the Medical Profession for a quarter of a century in the treatment of Dermatic Diseases, for cleansing fetid discharges, for making gynecological examinations, etc.

AS A SHAMPOOING AGENT PACKER'S TAR SOAP is undoubtedly the best known. It is extensively used, both as a delightful cleanser to preserve healthful conditions of the hair and scalp, and in treatment of dandruff and baldness.

THE PACKER MANUFACTURING CO., 81 Fulton Street. New York.

HENRY'S Three Chlorides

(LIQ. FERRISENIC HENRY'S).

THE GLANDULAR STIMULANT AND
CHALYBEATE TONIC.

Tonic, stomache, antiscorbutic,
hæmatinic and alterative.

In Anæmias, Chlorosis. A reconstructive in convalescence from any cause.

Makes Quantity and Quality of Blood.

DOSE: 1 or 2 drachms.

12 oz., \$1.00.

LITERATURE ON APPLICATION.

HENRY'S TRI-IODIDES.

(LIQ. SALT-IODIDE HENRY'S).

ALTERATIVE { Antirheumatic.
Antisyphilitic.

In lymphatic and visceral disturbances a uric acid solvent, and a reliable eliminant in all diathetic diseases.

8 oz., \$1.00.

1 or 2 fluidrachms in water exercises a synergistic effect over the entire range of

THE SALICYLATES AND IODIDES.

MAIZO- LITHIUM

(LIQ. LITHIUM MAIZENATE).

A POSITIVE DIURETIC.

STIMULATING, YET NON-IRRITATING.

In genito-urinary diseases, cystitis, incipient diabetes, lithæmia, urethritis, and all inflamed conditions requiring a non-irritating diuretic.

DOSE: 1 or 2 drachms.

8 oz., \$1.00.

HENRY PHARMACAL CO., LOUISVILLE, KY.

BENSOLYPTUS

Schieffelin's

Alkaline Antiseptic and Prophylactic.

Bensolyptus is an agreeable alkaline solution of various highly approved antiseptics, so combined that one supplements the action of the other. Its ingredients are all of recognized value in the treatment of

Catarrhal Affections

because of their cleansing, soothing, and healing properties. Bensolyptus is highly recommended in all diseases of the nose and throat, both acute and chronic, and as a mouth-wash and dentifrice. It is also indicated for internal use in affections of the stomach and intestines where an agreeable, unirritating, and efficient antiseptic and antifermentative is required.

Pamphlets on application to

Schieffelin & Co., New York.

Schieffelin & Co.'s

(NEW YORK)

Soluble Pills and Granules.

Reliable. Soluble. Beautiful.

The Perfection of Pill Making.

Examine Them and Notice

Their Perfect Uniformity in size and weight.
The Crystal Transparency of the coating.
They are the only pills which show the precise colors of the masses.

Test Them and Prove

Their Ready Solubility.
Their Exact Conformity to their several Formulas.
The Certainty of Proper Therapeutic Effects.

Our list of Pills and Granules embraces those made according to the formulas of the U. S. Pharmacopœia; also most of those in common use among the profession.

We also manufacture pills to order.

SPECIFY Schieffelin's WHEN PRESCRIBING.

In writing to advertisers mention Monthly Cyclopædia

ALWAYS THE SAME. A STANDARD OF ANTISEPTIC WORTH.

Listerine.

LISTERINE is a non-toxic, non-irritating and non-escharotic antiseptic, composed of ozoniferous essences, vegetable antiseptics and benzo-boracic acid.

LISTERINE is to make and maintain surgical cleanliness in the antiseptic and prophylactic treatment and care of all parts of the human body.

LISTERINE is invaluable in obstetrics and gynecology as a general cleansing, prophylactic, or antiseptic agent.

LISTERINE is useful in the treatment of the infectious maladies which are attended by inflammation of accessible surfaces—as diphtheria, scarlet fever and pertussis.

LISTERINE diluted with water or glycerine speedily relieves certain fermentative forms of indigestion.

LISTERINE is indispensable for the preservation of the teeth, and for maintaining the mucous membrane of the mouth in a healthy condition.

LISTERINE is of accurately determined and uniform antiseptic power, and of positive originality.

LISTERINE is kept in stock by all worthy pharmacists everywhere.

Lambert's Lithiated Hydrangea.

A valuable renal alterative and antilithic agent of marked service in the treatment of Cystitis, Gout, Rheumatism, and diseases of the Uric Diathesis generally.

LITERATURE UPON APPLICATION.

LAMBERT PHARMACAL CO., ST. LOUIS.

HIGHLAND WATER



*Is the Best Tonic, Diuretic, and
Solvent Eliminator of Uric Acid:*

1. **Because** it is the softest and purest water known, containing no excess of mineral salts and no lime or organic impurities, being therefore non-irritating to the digestive organs and kidneys.
2. **Because** it contains Iron in combination with the alkalies, Sodium and Potassium in small quantities, as recommended by Professors Flint, Thomson, Draper, and others.
3. **Because** it comes from a region free from malarial germs.
4. **Because** all its mineral salts, and five of them in the exact form given in its analysis, are found in the human blood, of which they form a most essential part.

At Grocers and Druggists, or

Office, 439 FIFTH AVENUE,

Telephone, 1763-38th St.

NEW YORK.

In writing to advertisers mention Monthly Cyclopædia

SVAPNIA

OR

PURIFIED OPIUM

FOR PHYSICIANS USE ONLY.

Contains the Anodyne and Soporific Alkaloids, Codeia, Narceia and Morphia. Excludes the Poisonous and Convulsive Alkaloids, Thebaine, Narcotine and Papaverine.

SVAPNIA has been in steadily increasing use for over twenty years, and whenever used has given great satisfaction.

To PHYSICIANS OF REPUTE, not already acquainted with its merits, samples will be mailed on application.

SVAPNIA is made to conform to a uniform standard of Opium of Ten per cent. Morphia strength.

JOHN FARR, Manufacturing Chemist, New York.

C. N. CRITTENTON, Gen'l Agent, 115 Fulton St., N. Y.

To whom all orders for samples must be addressed

SVAPNIA IS FOR SALE BY DRUGGISTS GENERALLY.

Continue to ask for Parke, Davis & Co.'s Anti-Diphtheritic Serum. We continue to lead in potency, efficiency and reliability of product. Remember also that P., D. & Co.'s serum is marketed in hermetically sealed bulbs (not ordinary vials and corks), which exclude the air and keep the serum strictly aseptic.

Latest literature mailed upon request.

Special Notice.—The validity of the recent patent issued by the U. S. Patent Office on diphtheria anti-toxin will be contested in the courts. Pending a decision in the case, we beg to announce that we shall, at our own expense, protect and defend physicians in any legal proceedings which may be brought as a result of the purchase of "P., D. & Co.'s" Anti-Diphtheritic Serum. Order it at any time and in any quantity without hesitation.

Parke, Davis & Co.,

Home Offices and Laboratories, Detroit, Mich.

Branches in New York, Kansas City, Baltimore, and New Orleans.

DOCTOR!
Antikamnia
 —> IS NEITHER <—
FRENCH NOR GERMAN
IT IS AMERICAN
MADE IN AMERICA
FOR AMERICANS

SYRUPUS ROBORANS.

Syr. Hypophos. Comp. with Quin., Mangan., and Strych.
 1-128 gr. Strychnine to teaspoonful.

Syrupus Roborans as a Tonic during Convalescence has no equal.

As a nerve stimulant and restorative in wasting and debilitating diseases, as a constructive agent in Insomnia, Pneumonia, Tuberculosis, Bronchial Asthma, Marasmus, Strumous Diseases, and General Debility, this compound has no superior.

SYRUPUS ROBORANS is in perfect solution and will keep in any climate.

Dr. T. H. Stucky writes: In a case of Tertiary Syphilis, very anæmic, the Iodides were revolting to the stomach, being vomited when taken. Syrupus Roborans given three weeks with improvement, when the Iodide of Potassium was retained with good results.

Dr. W. O. Roberts says: In cases convalescing from "La Grippe" Syrupus Roborans has no equal.

For Sale by all Wholesale Druggists.

PETER'S PEPTIC ESSENCE CO.

A Powerful Digestive Fluid in Palatable Form.

Please note that Essence and Elixir Pepsin contain only Pepsin, while in Peter's Peptic Essence we have all the digestive ferments. These are preserved in solution with C. P. Glycerin in a manner retaining their full therapeutic value, which is exerted in and beyond the stomach.

It is a Stomachic Tonic, and relieves Indigestion, Flatulency, and has the remarkable property of arresting Vomiting during pregnancy. It is a remedy of great value in Gastralgia, Enteralgia, Cholera Infantum, and Intestinal derangements, especially those of an Inflammatory character. For nursing mothers and teething children it has no superior. Samples sent upon application. Express charges at your expense.

ARTHUR PETER & CO., Louisville, Ky.

EUQUININE....



Tasteless, Nontoxic QUININE

ESPECIALLY INDICATED IN **CHILDREN'S CASES.**

A SPECIFIC AGAINST **WHOOPIING-COUGH.**

VEREINIGTE CHININLABRIKEN....

ZIMMER & CO.,

Frankfort O. M., - - Germany.

...VALIDOL

An Analeptic and Antihysteria

SPECIFIC AGAINST SEASICKNESS

R. W. PHAIR & CO., Agents, 16 Platt Street, NEW YORK.

In writing to advertisers mention Monthly Cyclopædia

HEMORRHOIDS PALLIATIVE TREATMENT

When internal Hemorrhoids have attracted the attention of the patient, they have usually progressed further in development than external Hemorrhoids and the physician is called upon to employ mild palliative or curative treatment, or radical operative procedures according to the requirements of any general case.

Palliative treatment in early cases sometimes becomes curative treatment. Dr. Carpenter reports some fifty cases of piles treated with Glyco-Thymoline (Kress), diluted with an equal portion of water. Two teaspoonfuls or more of this mixture are injected into the rectum with a small glass or hard rubber syringe, two or three times a day. The results obtained from this treatment were invariably satisfactory, giving immediate relief to the patient.

Glyco-Thymoline (Kress) causes rapid exosmosis of serum from the engorged tissues; it is perfectly bland and devoid of irritating qualities and being antiseptic it is particularly advantageous in cases in which abrasion or erosion of mucosa exists, making the employment of an antiseptic desirable.

One or two tablespoonfuls of Glyco-Thymoline (Kress) diluted one-half with water injected into the rectum at a time when the patient wishes to move the bowels, will give beneficial results.

GLYCO-THYMOLINE

(KRESS)

at once stops the bleeding, allays the pain, itching and discomfort.

KRESS & OWEN COMPANY

Manufacturing Chemists

MANUFACTURERS OF BIRMINGHAM NASAL DOUCHE,
MARKASOL, BISMUTH BORO-PHENATE.

221 FULTON ST., NEW YORK



IF THERE IS ANY DOUBT IN YOUR MIND AS TO THE SUPERIORITY OF....

WELCH'S GRAPE JUICE

over other grape products and malt preparations--send 10c. in stamps to us for a 3oz. bottle, or buy a pint bottle from your druggist.

WELCH'S GRAPE JUICE is Concord Grapes
---in Liquid Form

IT MAY BE GIVEN AS SOLE DIET.

In Wasting Diseases Welch's Grape Juice affords an immediate vigor from which there is no reaction, being readily assimilated by the blood. It stays down when nothing else will.

A Pint Bottle sent Free to any Physician who will Pay Express Charges.
Booklet sent on application

WELCH GRAPE JUICE CO., Westfield, N. Y.

In writing to advertisers mention Monthly Cyclopædia

ZE=AN (N., B. & Co.)

A Concentrated Fluid Extract of
Corn Silk.

Ze-an (N., B. & Co.) is the most active diuretic of the materia medica, a solvent of uric acid and phosphatic gravel and an anodyne to the genito-urinary mucous membrane.

Ze-an (N. B. & Co.) is *par excellence* the remedy for

**NEPHRITIS
CYSTITIS**

URETHRITIS
and for **URIC ACID** and
**PHOSPHATIC
GRAVEL.**

Useful also for the removal of dropsical effusions.

UTEROTONIC

(N., B. & Co.)

A combination in palatable form of Squaw Vine, Black Haw, Jamaica Dogwood, Golden Seal, Black Cohosh, Cascara Sagrada, Hyoscyamus and Potassium Bromide.

UTEROTONIC (N., B. & Co.) is the most successful remedy yet offered for atonicity and irritability of the uterus and ovaries,

Dysmenorrhœa,

Menorrhagia,

Ovarian Neuralgia,

Endometritis,

and as a preparation for **Parturition,**

Increasing the force and lessening the pain of the uterine contractions.

Full Descriptive Pamphlets and Samples of the above furnished on application to

NELSON, BAKER & CO.,

Manufacturing Pharmacists,

- DETROIT, MICH.



BAIRD STEAM FOMENTER

A UNIVERSAL MEDICAL
AND HOUSEHOLD APPARATUS

It consists of an alcohol heater, in connection with a boiler filled with water, to which a coil of rubber tubing is attached. When in operation, steam circulates through the coil, thus giving a hot fomentation (moist or dry) of even temperature.

The FOMENTER supplies the most efficient, economical and satisfactory method of applying both dry and moist hot fomentation ever offered the medical profession. It gives a continuous hot fomentation of even temperature. The degree of heat can be regulated to a nicety, and, if desired, the fomentation may be continued indefinitely. Patients need not be disturbed from their sleep and rest, nor exposed to sudden changes of temperature while changing the hot fomentation, as in the case of old methods.

In cases of Muscular Stiffness and Joint Diseases, the affected parts are wrapped in flannel with straight tubing wrapped around this. Fomentation may be continued at the degree of heat desired. Moist fomentation is produced by keeping the flannels moist.

The FOMENTER is also the best medium for producing medicated fumes for the treatment of all catarrhal affections of the head, throat and lungs. The medicine is placed in the water in the boiler. The warm inhalation causes the treatment to penetrate and heal the diseased parts, where the old method has no effect.

Canopies can be made of the bed blankets for use in cases of croup and diphtheria, and for vapor baths, etc. If the patient is confined to bed, steam can be conveyed under the blanket by means of a short piece of tubing. Diaphoresis can be continued as long as desired.

There should be one in every home and physicians should prescribe it to their patients.

The BAIRD STEAM FOMENTER is also valuable as a sterilizer for small instruments.

Particulars furnished on application.

Price, \$1.50 to \$3.50

BAIRD BROS. & CO., No. 170 Madison St., Chicago, Ill.

In writing to advertisers mention Monthly Cyclopædia

Guaiacol Carbonate

(Duotal.)

Creosote Carbonate

(Creosotal.)

These two Carbonates are now acknowledged to be the most pleasant, safe and effective remedies in the treatment of tuberculosis. An experience of over five years has established the fact that Creosote Carbonate and Guaiacol Carbonate are free from the noxious by-effects of the plain creosote and guaiacol, and exhibit only the therapeutic properties of these drugs. The neutral Carbonates cause neither irritation nor inflammation of the stomach or intestines, and abnormal fermentation and decomposition processes are stopped. No other guaiacol compound contains so high a percentage of chemically pure crystalline guaiacol as Duotal, namely: over 90%; while Creosotal contains 92% of pure creosote. The Carbonates are the only two intestinal antiseptics whose action extend as far as the ileum; but, as has been shown by thorough investigation, they cannot be found at the commencement of the large intestine when administered in ordinary doses.

Professor Dujardin-Beaumetz, in the Supplement to the Dictionary of Therapeutics, Paris, 1895, states:

"From numerous observations, published both in France and abroad, it appears that Carbonate of Guaiacol stimulates the appetite, facilitates assimilation, invariably produces a considerable increase in weight, and imparts to the system great resisting power. Its antiseptic action will arrest the development of any microbes in the intestine and obviate the generation of gases, which is always unpleasant. Fits of coughing will decrease, the expectoration will be modified, fever as well as perspiration at night will gradually disappear, and the patient will thus enjoy refreshing sleep."

Dr. Nied, Physician-in-Chief at St. Elizabeth's Hospital in Vienna, has treated last winter a number of female patients who were suffering from acute bronchitis, complicated in some cases by influenza, with Guaiacol Carbonate in doses of from $7\frac{1}{2}$ grains to $22\frac{1}{2}$ grains, twice daily, with excellent results. (*Allgemeine Wiener Medicinische Zeitung*, June, 1897).

Dr. Eschle in Professor Baumann's physiologico-chemical laboratory in the Medical Department of the University of Freiburg, found that Guaiacol Carbonate is the more thoroughly used up the smaller and more frequent the doses in which it is given. In phthisis, therefore, Eschle recommends the administration of a dose of $4\frac{1}{2}$ to $7\frac{1}{2}$ grains of Guaiacol Carbonate four times daily. (*Zeitschrift für Klinische Medizin*, Volume 29, 3 and 4).

Dr. Jacob, Chief at the University Medical Clinic of Professor Leyden in Berlin, has obtained very satisfactory effects from the use of Creosotal in a number of cases of pronounced phthisis, though comparatively small doses have been employed. Each patient began with 5 drops three times daily, increasing the dose three drops every day, until 25 drops were taken at a dose. At this they were kept for from one to four weeks, when the dose was diminished in a similar ratio, until only 10 drops were taken thrice daily; and then the ascending scale was begun again. (Inaugural Dissertation by Hans Nordt, University of Berlin, February, 1897.)

VON HEYDEN'S

ANTISEPTIC CREDE (Citrate of Silver, Heyden)
 CREDE'S LACTATE OF SILVER
 APOLYSIN
 GUAIACOL CHEMICALLY PURE CRYSTALS
 GUAIACOL PURE, LIQUID
 GUAIACOL-SALOL
 OLEO-CREOSOTE

OLEO-GUAIACOL
 ORPHOL (Dotanaphthol-Bismuth)
 PHENOL-DISMUTH
 XEROFORM (Tribromphenol-Bismuth)
 ORTHO-CHLORPHENOL
 PARA-CHLORPHENOL
 PARA-CHLORSALOL

SCHERING & GLATZ, 55 Maiden Lane, New York,
 Sole Agents for the United States and Canada.

Literature furnished on application.

In writing to advertisers mention Monthly Cyclopædia

DO NOT BE DECEIVED.

- DO NOT BE DECEIVED—*Maltzyme* is not an *imitation* of anything
- DO NOT BE DECEIVED—Chemistry and pharmacy *advance* like other departments of science, but malt products have not been improved—until *Maltzyme* was produced.
- DO NOT BE DECEIVED—*Maltzyme* represents improvement, *perfection* in diastasic malt products.
- DO NOT BE DECEIVED—*Maltzyme* is an *essence* of malt and the best diastasic product known.
- DO NOT BE DECEIVED—*Maltzyme* only entered the therapeutic field in October last, but it already *leads* all other products of its class right here at home in New York City.
- DO NOT BE DECEIVED—*Maltzyme* (plain) is the best preparation known for the treatment of starchy indigestion, being more diastasic than any other malt product, rich in nutriment, and is *not* thick and gummy.
- DO NOT BE DECEIVED—*Maltzyme* with Cod Liver Oil. No other preparation can hold a candle to it. Taste it. Try it. It has *twice* as much diastase as any other combination of oil and malt, and it is palatable.
- DO NOT BE DECEIVED—*Maltzyme* with Cascara Sagrada has in it all the virtues of the bark, more diastase than any other Cascara product, is limpid, and is *readily* taken.
- DO NOT BE DECEIVED—*Maltzyme* with Hypophosphites is not an *antiquated* formula. It contains Manganese, as well as Hypophosphites of lime, iron and potassium.
- DO NOT BE DECEIVED—The falsehoods told by jealous rivals amuse *you* very much, so we hear. They do us, certainly.
- DO NOT BE DECEIVED—Shall we send you our circular, and discuss the question of *samples* with you?

MALT DIASTASE CO., 1 Madison Ave., New York

Beer as a Remedy

Typhoid Fever

During convalescence from fever, when the heart and stomach have been weakened and the digestion is weak, malt liquors properly brewed increase the appetite, digestion and nutrition. (VARIOUS TEXT BOOKS ON THERAPEUTICS.)

American Physicians have at their disposal the best beer in existence

Tannhæuser and Old Stock Lager

THE WORLD'S JUDGMENT OF OUR PRODUCT

Philadelphia, 1876, Two Medals; Paris, 1878, Grand Prize; Brussels, 1888, Grand Gold Medal and Diploma of Honor; Paris, 1889, Grand Prize; Chicago, 1893, Four Prizes; Antwerp, 1894, Grand Prize; Amsterdam, 1895, Grand Prize; Brussels, 1897, Grand Prize.

The Bergner & Engel Brewing Co., Philadelphia

In writing to advertisers mention Monthly Cyclopædia

Phenālgin

NON TOXIC, HYPNOTIC,
Analgesic, Antipyretic,
NON DEPRESSANT, STIMULANT.

FORMULA

Ammonio-Phenylacetamide, pure white.
Made from C. P. material.
A stable and uniform product.

R Phenalgin (Etna), gr. 10.
Make one powder for
painful menstruation.

R Phenalgin (Etna), gr. 5.
Make one powder for
insomnia.

Generous Samples of
Powder or Tablets of
Pure Phenalgin Mailed
Free to Physicians.

ETNA CHEMICAL CO.

313 WEST STREET.

NEW YORK, U.S.A.

S GLASS & METAL S
HIGHEST GRADE
FOR PHYSICIANS,
DENTISTS ETC.
J.W. RUSSELL & CO.
MFRS. 294 PEARL ST. NEW YORK.
WRITE FOR PRICE LIST.

Send for the interesting and beautiful book entitled
**FACTS AND SUGGESTIONS REGARDING THE PI-
NELLAS PENINSULA, FLORIDA.** It is Illustrated
with 75 Beautiful Half-Tone Engravings, nearly all
of them from Photographs. Handsomely Printed on
Fine Enameled Paper. Crown Octavo. 132 Pages.
Price, 25 cents, postage prepaid. THE F. A. DAVIS
Co., Publishers, 1914-16 Cherry St., Phila., Pa.

PROGRESS IN SURGERY

has been largely due to antiseptic preparations. One great want
of the surgeon and general practitioner has been a surgical dress-
ing that was thoroughly antiseptic and applicable wherever
inflammation was present.

UNGUENTINE

fills this want. It is made upon the modified formula of Sir
Astley Cooper, the most renowned surgeon in the world in his
time. Its ingredients are Alum, Carbolic Acid, and Ichthyol,
combined with a pure Petrolatum base.

**ITS SUPERIORITY IS DUE TO THE FACT THAT
ALL IRRITATING PROPERTIES ARE ELIMI-
NATED FROM THE ALUM.**

This gives a surgical dressing of marvelous healing quali-
ties—that is astringent, but not irritating. It gives us a dressing
that is indicated wherever **inflammation** is present. The physi-
cian or surgeon who will give Unguentine a fair trial will find it the **best surgical dressing** ever made, with
the **widest field of usefulness.** It is convenient because always ready when wanted. Keeps in any climate.
Without acidity. Never grows rancid. **Heals without leaving scar tissue.** We ask the profession to give
Unguentine a fair trial, and judge it by what it does—not by what is said about it. Write to-day for a
free sample, clinical reports, and a biography of Sir Astley Cooper, originator of the working formula.



SIR ASTLEY COOPER.

THE NORWICH PHARMACAL CO.,

New York Salesroom, 51 John St.

NORWICH, NEW YORK.

AT DRUGGISTS' IN FOUR-OUNCE AND ONE-POUND JARS.

In writing to advertisers mention Monthly Cyclopædia

Maltine

MALTINE is not merely "malt," nor is it a mere "extract of malt," nor an "essence of malt."

MALTINE is the most highly concentrated extraction of all the nutritive and digestive properties of Wheat, Oats, and Malted Barley.

It has stood alone and unrivalled throughout the world in the therapeutic field for more than twenty years, despite the most strenuous efforts of the ablest pharmaceutical manufacturers to produce a preparation approaching it in medicinal value, elegance, palatability, and stability.

"Malt" is not "MALTINE."

"Extract of Malt" is not "MALTINE."

"Essence of Malt" is not "MALTINE."

"MALTINE" must be designated to get "MALTINE."

DOCTOR Do You Cure Your Goitre Patients?

I cure all of mine and receive from \$25 to \$50 from each of them. The cost of treating a case will not exceed \$1.00. Enclose a 2-cent stamp for particulars.

Lock Box 604.

F. E. MAY, M. D., Bloomington, Ill.

JOHN WELSH DULLES
EDWARD W. CRITTENDEN
DAVID SWOPE

Telephone 5149

DULLES & CRITTENDEN,

INSURANCE,

NO. 312 WALNUT STREET,
PHILADELPHIA.

FIRE
LIABILITY
BOILER
PLATE GLASS

PLANTEN'S
CAPSULES
AND **PERLOIDS**
OF
SANDAL OIL

CAPSULES.

No.	Size.	12 in Box	PER DOZ.
No. 53A	10 Minims	24	\$2.25
" 53	"	36	4.25
" 54	"	100	6.25
" 54C	"		15.00

"PERLOIDS,"

OR IMPROVED FRENCH "PERLES."
Cheaper and Better than the Imported.

No.	Size.	40 in Vial	PER DOZ.
No. 421A	5 Minims	80	\$4.75
" 421B	"	100	9.00
" 421C	"		10.20

A trial bottle or dozen sent prepaid on receipt of price.
H. PLANTEN & SON (Established), New York.
1836

Manufacturers of over 400 varieties of
FILLED AND EMPTY CAPSULES.

In writing to advertisers mention Monthly Cyclopædia

**Uniformly Effective, Agreeable and Lasting,—the
Standard Preparation of Erythroxylon Coca**

During past 30 years
most popularly used
Tonic-Stimulant in
Hospitals, Public and
Religious Institutions
everywhere.



We have received
over 7000 written
endorsements from
PROMINENT PHYSI-
CIANS in Europe and
America.

"MARIANI WINE"

FORMULA : The concentrated extract—the aromatic principle of the fresh Coca Leaf,
blended with a special quality of grape juice of southern France.

DOSE : Wine-glassful three times a day, or more or less at Physician's discretion.

Nourishes = Fortifies = Refreshes
AIDS DIGESTION - STRENGTHENS THE SYSTEM

AGREEABLE TONIC-STIMULANT WITHOUT UNPLEASANT REACTION.

To avoid disappointment please specify "Vin Mariani."

SOLD AT ALL PHARMACIES.

PARIS: 41 Boulevard Haussmann.
LONDON: 239 Oxford Street.
MONTREAL: 28 Hospital Street.

MARIANI & CO., 52 W. 15th St., New York.

AXION ELASTIC TRUSS.

WORN NIGHT AND DAY WITH COMFORT. HOLDS THE
WORST RUPTURE WITH EASE.



We are now manufacturing a superior grade Silk Elastic Webbing. This
Webbing, made of selected rubber, wound with Silk, has a strength, durability, and
softness of texture unequalled. This Webbing is used only in our **AXION ELASTIC TRUSS.**

Our **New Axion Pad** has now been in successful operation for six months, and is giving universal
satisfaction. It is the most durable and comfortable soft pad made. It can be made **firm** or **soft** at any
time by simply turning a screw. Perfectly adjustable in every way—it is adapted to every variety of
Hernia.

The **AXION ELASTIC TRUSS** holds rupture perfectly, exercising and resting, and is worn
with the greatest comfort by the patient.

SEND FOR ILLUSTRATED CATALOGUE.

G. V. HOUSE MFG. CO.,

744 Broadway, New York City.

Send for our New 28-Page Descriptive Catalogue. Mailed to any
address promptly on request. THE F. A. DAVIS COMPANY,
Medical Publishers, 1914-16 Cherry Street, Philadelphia, Pa.

The American
THERAPIST

A Monthly Record of Modern
Therapeutics.

Original, Practical,
Independent, Interesting.

SUBSCRIPTIONS SOLICITED.

THE AMERICAN THERAPIST PUBLISHING CO.,
Lock Box 1170, New York.

\$1.00
PER YEAR

Sample Copies, 10 cts.

In writing to advertisers mention Monthly Cyclopædia

CHIONIA

THE HEPATIC STIMULANT

INDICATED IN

Diseases Caused by Hepatic Torpor.

Does not purge, per se, but under its use the Liver and Bowels
gradually resume their normal functions.

DOSE—ONE TO TWO FLUID DRACHMS, THREE TIMES A DAY.

PEACOCK'S BROMIDES

THE STANDARD SEDATIVE

INDICATED IN

Congestive, Convulsive and Reflex Neuroses.

Absolutely uniform in purity and therapeutic power, produces clinical results
which can not be obtained from the use of commercial bromide substitutes.

DOSE—ONE TO TWO FLUID DRACHMS IN WATER, THREE TIMES PER DAY.

PEACOCK CHEMICAL COMPANY, St. Louis, Mo.

—AND—

36 BASINGHALL ST., LONDON, ENGLAND.

SENG

FOR

**INDIGESTION, MALNUTRITION, PHTHISIS,
AND ALL WASTING DISEASES.**

DOSE—One or more teaspoonfuls three times a day. For babies, ten to fifteen
drops during each feeding.

CACTINA PILLETS

FOR **ABNORMAL HEART ACTION.**

DOSE—One Pillet every hour, or less often as indicated.

SULTAN DRUG CO., St. Louis and London.

In writing to advertisers mention Monthly Cyclopædia

THE GREAT FACT IN MODERN MEDICINE:

"The Blood is the Life,"

*And Where Nature fails to make Good Blood,
WE CAN INTRODUCE IT.*

BOVININE is Bovine Blood Unaltered from the Arteries of the Bullock;
The Universal Auxiliary of Modern Medicine and Surgery,
and the TRUE "ANTITOXIN" of Healthy Nature.

In the more enlightened progress of Modern Medicine, "Blood-letting" has given place to Blood-getting.

Aye! Get Good Blood—but How? Not by the Alimentary Process. It has already failed to do its work (else the patient would not be sick); and in acute disease must not even be allowed to do the work it can. Stimulate as you will, the whole sum of the patient's alimentary power when fully forced into play, is unable to keep up the nourishing and supporting contents of the blood. There is absolutely but one thing to do; and, thank God, that can be done, usually with success, as ten-thousand-fold experience has proved. That one thing is this: where Nature fails to PRODUCE good and sufficient Blood, WE CAN INTRODUCE IT from the arteries of the sturdy bullock, by the medium of BOVININE.

The vital activity of this living blood conserve rests on no man's assertion: it speaks for itself, to every properly equipped physician who will test its properties microscopically, physically, or therapeutically.

TRY IT IN PRACTICE.

TRY it in Anæmia, measuring the increase of red cells and hæmaglobin in the blood as you proceed, together with the improving strength and functions of your patient.

Try it in Consumption, with the same tests from week to week.

Try it in Dyspepsia or Malnutrition of young or old, and watch the recuperation of the paralysed alimentary powers.

Try it in Intestinal or gastric irritation, inflammation, or ulceration, that inhibits food itself, and witness the nourishing, supporting and healing work done entirely by absorption, without the slightest functional labor or irritation; even in the most delicate and critical conditions, such as Typhoid Fever and other dangerous gastro-intestinal diseases, Cholera Infantum, Marasmus, Diarrhœa, Dysentery, etc.

Try it per rectum, when the stomach is entirely unavailable or inadequate.

Try it by subcutaneous injection, when collapse calls for instantaneous blood supply—so much better than blood-dilution!

Try it on Chronic Ulceration, in connection with your antiseptic and stimulating treatment (which affords no nourishment) and prove the certainty and power of topical blood nutrition, abolishing pus, stench, and PAIN, and healing with magical rapidity and finality.

Try it in Chronic Catarrhal Diseases; spraying it on the diseased surfaces, with immediate addition of peroxide of hydrogen; wash off instantly the decomposed exudation, scabs and dead tissue with antiseptic solution (Thiersch's); and then see how the mucous membrane stripped open and clean, will absorb nutrition, vitality and health from intermediate applications of pure bovinine.

Try it on the Diphtheritic Membrane itself, by the same process; so keeping the parts clean and unobstructed, washing away the poison, and meanwhile sustaining the strength independently of the impaired alimentary process and of exhaustive stimulants.

Try it on anything, except plethora or unreduced inflammation; but first take time to regulate the secretions and functions.

Try it on the patient tentatively at first, to see how much and how often, and in what medium, it will prove most acceptable—in water, milk, coffee, wine, grape, lemon or lime juice, broth, etc. A few cases may even have to begin by drops in crushed ice.

A New Hand-book of Hæmotherapy for 1898, epitomizing the clinical experience of the previous three or four years, from the extensive reports of Hospital and private practice. To be obtained of

THE BOVININE COMPANY, 75 W. Houston Street, New York.

LEEMING, MILES & Co., Montreal, Sole Agents for the Dominion of Canada.


In writing to advertisers mention Monthly Cyclopædia

Boil Our Syringe

AND THUS KEEP IT ASEPTIC. IT IS
SIMPLE, STRONG, COMPACT,
CLEAN.

\$3.00 In Handsome Aluminum Case

Containing Six Tubes of Our Soluble
Hypodermic Tablets.



DO NOT BOIL OUR HYPODERMICS

because it is unnecessary. They dissolve almost instantly in **cold** water. Those physicians who have been using tablets that dissolve slowly even in **hot** water will appreciate the superiority of "cold water" tablets, especially in emergency cases. Free samples.

SHARP & DOHME,

CHICAGO.


BALTIMORE.

NEW YORK.

The Best Medical Electrical Apparatus In the World.

TREATISE ON ELECTRO-THERAPY
FREE.

300 pages. 450 Illustrations.



Giving complete electrical technique of such operations as **Reduction of Strictures**; **Treatment of Dysmenorrhœa**; **Removal of Superfluous Hair, Warts, Moles, Nævi, &c.**; **Cataphoresis**; **Electro-Diagnosis**; **Metallic Electrolysis**; also giving prices and descriptions. We are desirous of placing this *educator* in the hands of every physician in the land, and will send it **post-paid, without charge**, if you will write and ask for it.

McINTOSH BATTERY AND OPTICAL CO.,

521-531 Wabash Avenue, CHICAGO, ILL.

In writing to advertisers mention Monthly Cyclopædia

IMPERIAL GRANUM

THE STANDARD PREPARED FOOD

IS the enviable possessor of a reputation
INCONTESTABLY ESTABLISHED
 in THOUSANDS OF CASES where it was THE ONLY NOURISH-
 MENT the stomach would tolerate, when LIFE SEEMED DEPENDING
 on its retention. IT IS —

UNRIVALLED IN TYPHOID
 And all **FEVERS,**

and in all gastric and enteric diseases especially in
DYSENTERY, DIARRHOEA, CHOLERA-MORBUS and CHOLERA INFANTUM.

Literature and SAMPLES for clinical test, supplied ONLY to PHYSICIANS and trained nurses.
 Sent FREE, charges pre-paid, on request. CORRESPONDENCE SOLICITED.

JOHN CARLE & SONS, 153 Water Street, NEW YORK CITY.
 SOLD BY DRUGGISTS generally.

Tannopine, a specific in diarrhoeal affections.

Tannopine (hexamethylen-tetramine tannin) is always prompt and reliable in action, because it passes unchanged through the stomach and is gradually decomposed in the intestine, thus exerting its astringent effect at the **site of disease**, whether in the upper or lower intestinal canal. Especial indications: Gastro-enteritis, cholera infantum, typhoid, and intestinal tuberculosis.

SCIENTIFIC FOODS, TONICS AND RESTORATIVES.

SOMATOSE,	FERRO-SOMATOSE,	LACTO-SOMATOSE,
a scientific food in all conditions of lowered vitality, producing a rapid gain in flesh and strength without taxing the digestive organs.	aferruginous nutrient in all anæmic states, palatable, readily assimilated, does not affect the teeth nor produce constipation.	pre-eminently a food for gastro-intestinal diseases, agreeable, nutritious and exerting a mild astringent action in diarrhoeal affections.

Protargol an important advance in gonorrhoea therapeutics

Protargol is regarded by Professor Neisser as "the best, safest and most rapid remedy hitherto introduced in the treatment of gonorrhoea." It is also highly recommended in affections of the eye, infected wounds, and as a substitute for nitrate of silver in general.

Send for samples and literature to

FARBENFABRIKEN OF ELBERFELD CO., 40 Stone Street, New York.

Selling Agents for the Bayer Pharmaceutical Products:

Aristol, Euophen, Ferro-Somatose, Hemicranin, Iodothyrene, Lacto-Somatose, Losophan, Lycetol, Phenacetin, Piperazine-Bayer, Protargol, Salicylic Acid, Salophen, Somatose, Sulfonal, Tannigen, Tannopine, Trional.

In writing to advertisers mention Monthly Cyclopædia

Special attention of the Medical Profession, is called to Colden's Liquid Beef Tonic.

It is composed of pure Extract of Beef, (Liquid) this is prepared by a special process. It is combined with a small per cent (25 per cent.) of Brandy, Cit. of Iron, Gentian, Peruvian and Calisaya barks. Promoting tissue formation, it is indicated in adynamic states, e. g., long slow convalescence after fevers, Anaemia, Anorexia. Delicate mothers with nursing infants, and those in whom debilitated conditions exist, derive untold benefit. It is a valuable nutritive tonic and stimulant, and well worthy of the reputation it has gained.

Colden's Liquid Beef Tonic

can be relied upon, and a faithful trial of the Preparation will prove this fact, On the label designated No. 1 with Iron, and No. 2 without Iron. Dose one teaspoonful to two tablespoonfuls three times a day. Children one teaspoonful three times a day. Sold by druggists generally.

MANUFACTURED BY

The Charles N. Crittenton Co.,
Laboratory,
115-117 Fulton Street, New York.

THE MONTHLY CYCLOPÆDIA OF PRACTICAL MEDICINE.

Vol. XII.
Old Series.

PHILADELPHIA, SEPTEMBER, 1898.

Vol. I. No. 9.
New Series.

TABLE OF CONTENTS.

PAGE	PAGE	PAGE
ABDOMINAL SECTION. H. T. Byford..... 344	GASTRO-ENTERITIS..... 351	MYXEDEMA IN THE NEGRO..... 356
ADENOID VEGETATIONS AND DEAF- MUTISM. E. Fayette Smith..... 345	Treatment. Wells, Charlotte C. West 351	Symptoms. Berkley..... 356
ADENOIDS, POST-NASAL..... 345	GENU VALGUM. Clarke..... 351	Treatment. Berkley..... 356
Diagnosis. Rice..... 345	GERMAN MEASLES..... 352	NEURASTHENIA..... 356
ALGINATE OF IRON. William Mac- lennan..... 346	Diagnosis. Forchheimer..... 352	Etiology. Frederick Peterson..... 356
AMENORRHEA..... 321	GONORRHEA..... 352	OPHTHALMIA NEONATORUM. Lucien Howe..... 357
Etiology. Alexander Simpson, Robert A. Reid..... 321	Etiology. H. Brooks Wells..... 352	PEMPHIGUS..... 357
Treatment. Leuf, W. E. Fothergill, Alexander R. Simpson, C. Earle Williams, Lutaud..... 322	Treatment. G. K. Swinburne..... 352	Treatment. R. F. Izlar..... 358
ANIMAL EXTRACTS. Parry, W. E. Moseley..... 346	GOUT..... 324	STERILITY..... 331
BLENNORRAGIC ARTHRITIS..... 347	Diagnosis. Watson Williams, A. H. Buck..... 324	Etiology. G. Pujol, E. Rode, Vedeler, Benzler..... 331
Treatment. König..... 347	Etiology. Froelich, Nobecourt, Cor- nillon..... 324	Treatment. Jones, W. Gill Wylie..... 332
BOILS, CARBUNCLES, AND FELONS... 347	Pathology. N. S. Davis (Jr.), A. P. Luff, C. S. Bull, Oliver, Riehl, Kittel..... 325	YELLOW FEVER..... 333
Treatment. L. Duncan Bulkley..... 347	Treatment. Luff, Ransom, Arm- strong, H. C. Wood, George W. Tobias, F. Levison, von Noorden, Grawitz, R. Newman..... 328	Diagnosis. John Guitéras, S. E. Archinard and R. S. Woodson, R. D. Murray, Eugene Wasdin, H. R. Carter, Melier, Edwin Kleba... 333
BRIGHT'S DISEASE..... 347	HYDROCEPHALUS, ACQUIRED..... 353	Bacteriology. Havelburg, Sanarelli, Surgeon-General Sternberg, E. Klebs..... 334
Treatment. James Tyson..... 347	Etiology. Bruce and Stiles..... 353	Pathology. Cuban Commission of Mississippi, R. D. Murray..... 337
CHLOROSIS..... 348	Treatment. Bruce and Stiles..... 353	Prognosis. Geddings..... 338
Treatment. Rosin..... 348	HYSTERIA AND PELVIC DISEASE. F. X. Dereum..... 354	Treatment. Sanarelli, E. K. Sprague, H. M. Folkes, H. D. Geddings, R. D. Murray..... 338
COLOTOMY AND COLOSTOMY. König and Sonnenburg, Mosetig-Moorhof 348	INTESTINAL PERFORATION..... 354	Prophylaxis. H. R. Carter, W. F. Brunner, Wasdin, Seaton, Nor- man, J. H. White..... 340
CONDURANGO. Lemoine..... 349	Diagnosis. E. M. Sutton..... 354	Post-epidemic Disinfection. Surgeon- General Walter Wyman..... 342
CONVULSIONS, INFANTILE..... 349	KRYOPHINE. John H. Curtis..... 355	BOOKS AND MONOGRAPHS RE- CEIVED..... 348
Etiology. Meunier..... 349	MEASLES..... 355	EDITORIAL STAFF..... 360
DYSENTERY. W. S. Attygalle..... 349	Diagnosis. Slawyk..... 355	
ENTEROCLYSMS. Judson Daland..... 349	MILK AS A CULTURE-MEDIUM. J. W. Strickler..... 355	
EXOPHTHALMIC GOITRE..... 350	MILK: ITS ABSORPTION vs. ITS DI- GESTION. Bulkley..... 355	
Treatment. Bertram..... 350		
FALLOPIAN TUBES, DISEASES OF..... 350		
Diagnosis. William H. Skene, A. J. C. Skene..... 351		

Cyclopædia of the Year's Literature.

AMENORRHEA.

Etiology.—Alexander Simpson¹ argues that just as modern pathologists have suggested that in the case of anuria uræmia results, so in the case of amenor-

rhea, where the menstrual flow does not occur, or does not escape, there is an element retained in the blood which

¹ Practitioner, Aug., '98.

causes a general disturbance in the system that might be called "menorrhæmia." The nervous system is at fault in many cases of amenorrhœa. In some cases patients get a chill, and that, perhaps, first affects the nervous system, while in others there is simply a mental impression, in which the patient has either a dread of conception or an eager hope of conception, and it has been noted that under one or the other of these emotional states the menstrual flow will sometimes be arrested for a month or two. So also in cases where the patient has had a fright or a disappointment or has been subject to a fit of passion. For the healthy performance of menstruation it is absolutely necessary that there should be a healthy circulatory apparatus and a healthy circulating fluid. Here, again, we find that in many of the cases of amenorrhœa there are defects in the circulatory apparatus, such as heart disease, or there is imperfect development of the blood-vessels. Everything which depresses the general health in a female is apt to show itself in depressing the menstrual function, in some cases even to its complete disappearance. But besides looking to these necessities for menstruation,—a healthy nervous system, good blood, and a good circulatory apparatus,—one must look to the seat of the menstrual flow and the associated organs, because the menstrual function is dependent not only on the condition of the uterus, but also on the condition of the associated organs. There is a tendency to refer the menstrual flow merely to conditions of the uterus and to the nervous system and the blood; but we must keep very strictly in view that the ovaries play a most important part in determining the regular healthy occurrence of menstruation. It has recently been shown that the nerve-

influence of the ovary is serviceable to the uterus in that it keeps up the steady rhythmical contractions that go on in the unpregnated uterus. So where the ovaries have become diseased, or have had to be removed, the uterine contractions that should go on normally, and become more pronounced at the menstrual periods, are absent. In those cases where the follicular stroma of the ovary has been the seat of an inflammatory process during the infectious fevers, the patient may have an amenorrhœa which may remain and become permanent.

An interesting case has been observed by Robert A. Reid,² in a young woman who presented many of the usual signs of pregnancy, including cessation of the menses, prominence of the abdomen, etc. On examination enormous deposits of adipose tissue were found in the abdominal walls, while the uterus was small—smaller indeed, than usual. Subsequent events proved it to be a case in which obesity had led to disturbance—if not, indeed, to early disappearance—of the menstrual function.

Treatment.—Leuf³ states that amenorrhœa may be caused by an underdevelopment of the uterus, or its atrophy, or to some neurotic influence. Cessations due to wasting diseases are conservative, the flow simply stopping because nature cannot permit the loss of blood in view of the other strain upon the system. An atrophied uterus requires the same treatment as an underdeveloped one, though it is less likely to respond to stimulation if it has existed for a long time, for retrograde changes are generally not as easily reversed as natural development is stimulated. The flow is most readily stimulated by the bipolar application of

² Mass. Med. Jour., Aug., '98.

³ Med. Council, Oct., '97.

the long, fine-wire, secondary faradic coil two or three times a week for from two to six months, continuing for ten or fifteen minutes at each sitting. It has also been accomplished by the passage of a similar current through the pelvis from the sacrum to the hypogastrium, the anode being behind. When the flow is simply scanty, the galvanic current is better, the cathode being in the uterine cavity and the anode upon the abdomen, with a current-strength of from 50 to 75 milliamperes. The leading fact should not be forgotten that the cathode increases circulation and nutrition, and that the anode has the opposite effect.

W. E. Fothergill⁴ claims that *senecio* will not cause abortion, or in any way influence the course of pregnancy. The practical utility of a drug which will cure functional amenorrhœa, but will not interfere with pregnancy, is, of course, obvious. A girl, for instance, is brought to the practitioner by her mother, amenorrhœa being the only symptom complained of. A physical examination is, for the time, out of the question, and a series of interrogations may cause unnecessary suffering. Functional amenorrhœa may be the condition or pregnancy. Under these circumstances, *senecio* may be safely prescribed before making a diagnosis, as it will probably cure the one, and certainly will do no harm to the other.

Eleven cases observed of true functional amenorrhœa in which *senecio* gave excellent results. No general disease, such as anæmia, or phthisis, was present, nor was there any deficiency, congenital or other, of the reproductive organs. The nervous mechanism which initiates the menstrual flow was, however, inactive in each case, and *senecio* appeared to be effective in stimulating it into action. In cases of anæmia, however, and

other conditions of exhaustion, due to disease, *senecio* has been found quite inactive in restoring menstruation. In such cases the cause of amenorrhœa is, of course, that the patient has no blood to spare; and treatment by a direct emmenagogue cannot be expected to have any effect, while indirect treatment by food, iron, etc., is indicated.

According to Alexander R. Simpson,⁵ no practitioner wants to merely stimulate uterine flow unless he also builds up the patient's system. In most cases the iron, when prescribed medicinally, has to be administered for some length of time. In recent times the preparation that has most favor is the Bland preparation of iron, whether in the form of a pill, tabloid, or a palatinoid. These preparations are given for at least six weeks to an amenorrhœic girl with pale mucous membranes. Where the iron of itself does not produce improvement in the blood, it sometimes brings about menstruation if it be given in combination with some other minerals—such as preparations of zinc or arsenic. As a rule, arsenic and iron bring about the conditions favorable to menstruation. Manganese, also, is very valuable in the treatment of amenorrhœa. In the managements of these patients it is important to keep in view that many of them are the subjects of constipation; hence, aloes-and-iron pill is a very favorable preparation.

In a clinical report on the value of ferratin in anæmic conditions Dr. C. Earle Williams⁶ reports the following interesting case, typical of frequent occurrences in general practice:—

T. P., age 19, had not menstruated for

⁴ Edinburgh Med. Jour., May, '93.

⁵ Practitioner, Aug., '98.

⁶ Amer. Therapist, Aug., '97.

five months. Her complexion was chlorotic, with large, dark rings around the eyes. There was dyspnœa and palpitation on the slightest exertion, constipation sometimes lasting seven days. Blood-count showed: red cells, 3,000,000; hæmoglobin, 52 per cent. She was placed on ferratin, 8 grains four times daily, with alœtic purges, combined with perfect rest. The dose of the ferratin was gradually increased to 12 grains four times a day, resulting in complete recovery. All the distressing symptoms left; menstruation returned, quantity and quality good; red cells, 4,600,000; hæmoglobin, 92 per cent.

The following formula has been followed by good results in amenorrhœa:—

R Strychnine sulphate, $\frac{1}{2}$ grain.
Iron peptonate,
Manganese lactate,
Scammony, of each, 20 grains.

To be divided into forty pills. Two to four pills to be taken every night on going to bed. (Lutaud.⁷)

GOUT.

Diagnosis.—Watson Williams⁸ states that in some countries gouty affections of the throat are even more common than rheumatic affections. The symptoms are very similar, though the pain may be more intense. Small tophi have been seen on the vocal cords and at the crico-arytenoid joint, though this condition is exceedingly rare. Gouty deposits in the laryngeal mucous membrane have been diagnosed as cancer. When gout has been correctly diagnosed as the pathological cause of the throat disease, the only local treatment should be in the form of mild, soothing sprays. Internally, colchicum and its preparations, iodides, Vichy water, etc., with suitable diet and hygiene should be given.

A. H. Buck⁹ claims that a patch of eczema, of spontaneous development, signifies the existence, in the subject possessing it, of a gouty diathesis. As it happens, the floor of the external auditory canal is apt to be the very first spot on the surface of the body where an eczematous inflammation develops. This condition constitutes a valuable guide-post, pointing as it does, at a very early stage, to the existence of that disturbed state of the metabolic processes to which the term "goutiness," or a "gouty diathesis," is applied. Different localities or different tissues are involved at different times. At one time it is the skin of the external auditory canal and auricle; at another time the dermoid surface of the tympanic membrane, as well as the walls of the canal; at still another the adjacent bone tissue is affected; and, finally, the disease locates itself in the mucous membrane of the tympanic cavity, and particularly in that part of it which borders upon the two fenestræ.

Etiology.—Froelich¹⁰ remarks that the theory must be regarded as proved which attributed the formation of uric acid in the body to a process of leucolysis following on a leucocytosis. A review of the literature upon tissue-necrosis in gout leads the writer to state definitely that the process is caused by a poison, probably a nucleic acid, acting in a similar way to that in which tissue-necrosis is caused by lead in plumbism.

Nobecourt¹¹ says that a form of gout exists related to saturnine intoxication. It is slowly established. The patient does not exhibit the various morbid

⁷ Jour. de Méd. de Paris, Dec. 19, '97.

⁸ Laryngoscope, Apr., '98.

⁹ Amer. Jour. Med. Sci., Mar., '98.

¹⁰ Jour. Amer. Med. Assoc., Jan. 3, '97.

¹¹ Sem. Méd., Apr. 23, '97.

phenomena of latent gout common in subjects of this diathesis throughout their early life. He is subject rather to colic, arthralgia, encephalopathy, apoplexy, albuminuria, paralysis, or dorsal tumor of carpus. He is unlike the hereditarily gouty, pale, thin, and anæmic. The mode of evolution of saturnine gout is analogous to that of ordinary gout. It resembles in some respects the gout of patients debilitated by any cause. The distinguishing features are frequent repetition of the attack, rapid generalization of the arthritis, and one frequent solitary visceral manifestation, namely: interstitial nephritis. Less-common phenomena are the rapid transformation of the acute into the chronic form, and the early development of tophi. Diagnosis is based on the same considerations as in ordinary gout. Again, gout in a saturnine patient is not necessarily saturnine gout.

Cornillon¹² remarks that, although the primary manifestations of gout are rarely initiated by traumatism, an acute attack of gout already pre-existing may be thus begun. The writer, however, details cases in which injuries were followed by the first appearance of the disease. Thus, a man of 53 years fell, dislocating his left shoulder. The dislocation was reduced at once, but three days afterward the left knee became swelled, red, and hot. The next day the great toe on the same side became similarly affected. Several days afterward all the joints were free of gout, but the right knee presented symptoms of it at a later date. The administration of salicylate of soda arrested the attack in about two weeks. This patient might probably have developed gout at some period or other, whether injured or not, but the injury precipitated the external symptoms of an inherent dyscrasia.

Pathology.—N. S. Davis, Jr.,¹³ states that the atheromatous changes which take place in the arterioles throughout the body, including those of the kidney, in the gouty subject are well known. But some points in connection therewith are often overlooked. One of these is the gradual progress of the renal disease, the organ being affected in spots, with intermissions in the degenerative changes which are microscopical in size, until finally large areas are involved. In these cases the glomeruli and tubules are attacked in a way at times to cause scarcely an appreciable symptomatology, whereas the same changes coming on suddenly, as in cases of a different etiology, cause striking clinical and urinary manifestations. The arterial changes in the nervous system lead to various nervous disturbances by interference with the nutrition of nerve-centres. Cerebral manifestations may arise from uræmia or from thrombosis of the cerebral arteries.

A. P. Luff¹⁴ believes that a functional affection of the kidneys always precedes gouty manifestations, and that this functional lesion, which may be started by various agents and causes (among which are excessive indulgence in nitrogenous foods, wines, and beer; the toxic effect of lead; and the influence of nervous impulses, such as mental shocks, severe accidents, etc.), may subside on the removal of the exciting cause, or it may pass on to a structural lesion, which is then of the contracted granular type and may also be transmitted by heredity.

The anatomical seat of the kidney affection is probably in the epithelium of the convoluted tubes, as that is the

¹² Progrès Méd., Jan. 2, '97.

¹³ Med. Rec., July 10, '97.

¹⁴ Indian Med. Rec., July 1, '98.

primary seat of disease in granular kidney and the increase of interstitial tissue is probably a secondary change, while granular kidney is not always evidenced by the occurrence either of albuminuria or of dropsy, and during life there may be no external manifestations of the existence of such renal mischief as the post-mortem may disclose.

On the whole, there is abundant evidence showing the direct connection between kidney trouble and gout, since uric acid has always been found in the blood in cases of renal disease, while uratic deposits are frequently found in the kidney and joints of gouty subjects and in the joints of persons with renal disease, but who have never been known to have had ostensible gout, and kidney mischief is frequently met with at the post-mortem of gouty subjects.

Alterations in the metabolism of the liver necessarily affect the formation, excretion, and daily elimination of uric acid by healthy persons; and, as variations in the metabolism of the liver are induced by changes in the quality and quantity of food ingested, by the amount of exercise, and by various nervous influences, it can be readily understood why liver trouble of some kind or another frequently accompanies gouty dyspepsia, and the fact is explained how several observers, unable to dissociate the connection between liver troubles and gout, have attributed the formation of uric acid to the liver.

The uric acid formed in the kidneys is at once converted into the quadriurates of ammonium, potassium, and sodium, and in healthy persons excreted dissolved in the urine, from which they sometimes separate, on cooling, as a deposit of "amorphous urates." In gouty conditions these urates are absorbed into the blood, where the sodic carbonate

converts the ammonium and potassium quadriurates into sodium quadriurate, which is an unstable salt and is gradually transformed into the less soluble and less easily excreted sodium biurate, which first passes into the hydrated or gelatinous modification, but, if it be present in the blood in greater amount than that fluid can retain in solution, it passes with the lapse of time and increasing accumulation into the almost-insoluble anhydrous or crystalline condition, and is deposited in those tissues (of the connective-tissue class) which, either on account of having received previous slight injuries or because of their poor vascular supply, favor its deposition.

C. S. Bull¹⁵ finds, associated with gout, changes in the walls of the blood-vessels of the retina, choroid, and optic nerve, including arteries, capillaries, and veins. Retinitis of a peculiarly-localized character, confined to the posterior zone of the fundus, with or without hæmorrhages in the retina and vitreous, and characterized by a peculiar yellowish exudation, occurring in clearly-defined patches. Optic neuritis, generally with, but sometimes without, an accompanying retinitis. The changes in the fundus are always bilateral, though rarely symmetrical in the two eyes. The lesion may begin simultaneously in the two eyes; but this is by no means always the case. The degenerative changes in the walls of the blood-vessels, both arteries and veins, are at first very minute and often overlooked. The general angiosclerosis and the patchy exudation in the retina cause marked impairment of central vision, but little impairment of the peripheral vision, and

¹⁵ Trans. Fourth Congress of Amer. Phys. and Surg., page 4, '97.

the disease never ends in blindness. The loss of central vision is always progressive up to a certain point. Improvement of the vision after the retinal disease is established cannot be expected, though in favorable cases the existing vision may be maintained. Hæmorrhages into the retina are rare except in the comparatively early stages of the disease. The most-marked feature in the ophthalmoscopic picture is the development of the angiosclerosis in the vessels of the retina.

Another almost equally marked symptom is the peculiar yellowish granular exudation in the retina, located by the ophthalmoscope around the posterior pole of the eye, and generally leaving the macula intact until late in the course of the disease. This exudation is shown by the microscope to be mainly in the nerve-fibre layer, though found in all the layers except that of the rods and cones. The changes in the optic nerves seem generally to be intra-ocular, but have been traced occasionally for some distance back of the eyeball.

Oliver¹⁶ says that the eyeball is nothing more than a lymph-holding end-organ, constantly changing its fluid constituents, and that in these structures changes can be produced by uric acid and urates. The presence of material in the outer coats of the eye that is so similar to that found in any joint shows how the external coverings of the organ and its encompassing capsule can be seriously disturbed by fluid change, inflammatory exudate, and inorganic deposit as the result of gouty diathesis. The lids, conjunctiva, and lacrymal apparatus are often swelled and excruciatingly painful. The inflammation is very rapid in onset, and quickly subsides by the application of dry heat and employment of constitutional measures.

Calcified Meibomian glands are frequently seen in middle-aged and elderly subjects as evidence of similar changes found elsewhere. Keratitis (peculiarly band-like in character) has been described by some authorities. The sclera gives evidence, particularly in the male subjects, by a series of extremely fugitive signs. Generally circumscribed, the redness is not rarely associated with intense dread of light and frequent attacks of severe pain, with copious lacrymation. The condition suddenly disappears by the local use of heat and the internal administration of large doses of alkalis. The iris is often also involved. In the ciliary body and choroid the diagnosis is less easy and the evil consequences are greater. Where the retina and optic nerve are concerned the ophthalmoscopic picture is generally typical. Hæmorrhages with changes in the size of the blood-vessels often appear early in the case. Later, shrinkage, atheroma, and sclerosis of the vessel-walls are apt to appear. At times these are associated with peculiar, glistening, yellowish bodies found in the region of the macula.

Riehl¹⁷ argues that in gout urate crystals are deposited in unaltered living tissues, and that the assumption that necrosis of the tissues must precede the deposit of the salts must be given up. Whether the tissue-increase about the urate deposits is due to mechanical or chemical irritation is not yet clear. Gouty nodules are probably formed as follows: First, a deposit of crystals in the lymph-spaces and vessels, and in the connective tissue itself, due to this inflammation of the surrounding tissue with granulation-tissue and giant-cells. The failure to find urates in the healthy

¹⁶ Jour. Amer. Med. Assoc., July 31, '97.

¹⁷ Wiener klin. Woch., No. 34, '97.

tissues is due, in part, to having examined tissues from the cadaver, and, in part, to faulty methods of hardening.

Kittel¹⁸ describes, in the aponeurosis of the foot and in the bones, a collection of small particles resembling concretions of sand, which may, by the pressure, interfere with the locomotion of the patients. In his opinion, these are the result of interference with the circulation of the foot, especially long-continued exposure of the foot to wet. Under these conditions a gradually-progressive degeneration takes place, which is followed by a necrosis of the tissues; a deposit of urates follows; stiffness, deformity, and distortion of certain of the joints takes place. The clinical symptoms are not those of true gout, but are considered by the writer to be a typical form of gout. Medicinal treatment against gout is useless, as the deposits of the urates are well encapsulated in necrotic tissue beyond the reach of any means of promoting absorption. For treatment the writer recommends mechanical exercise with massage, accompanied by cloths wet in alkaline solutions.

Treatment.—Luff¹⁹ says that the treatment of gout should aim at (1) checking excessive formation of uric acid in the kidneys, (2) preventing its absorption into the blood, and (3) promoting the removal of uratic deposits by facilitating the elimination of the quadriurate and biurates contained in the fluids of the body.

To check the excessive formation of uric acid, liver-metabolism should be promoted, and congestion of the portal system relieved by regulating the diet and regimen. Colchicum and guaiacum, as stimulants of hepatic metabolism, are very useful in many forms of gout. Constipation and the congestion of the

portal system may be relieved by occasional doses of blue pill followed by an Epsom-salts purge.

To promote the elimination of the quadriurates formed in the kidneys and so prevent their absorption into the blood is to strike at the primary evil in the causation of gout. To promote this diuresis should be increased and the activity of the urine diminished. Citrate of potassium is a good diuretic which not only increases the solubility of the quadriurates, but also diminishes the acidity of the urine and should be pushed until moderate alkalinity of the urine is produced.

The removal of uratic deposits and the elimination of quadriurates and biurates from the system may be attained by free diuresis, baths, and suitable exercise, and the careful selection of a mixed diet with a fair amount of vegetable food, since the mineral constituents of certain vegetables, such as Brussels sprouts, cabbage, French beans, spinach, turnips, and turnip tops possess to a remarkable degree the double function of inhibiting the conversion of sodium quadriurate into the biurates and increasing the solubility of the latter; but the idiosyncrasy of each patient to various articles of diet must be made the subject of careful observation.

Ransom²⁰ regards the use of drugs, other than tonics and digestives, in cases of gouty diathesis which show no actual symptoms of the disease as of doubtful value. The writer's treatment consists of six tumblerfuls of water daily, three of which may be taken hot; small doses of calomel followed by sulphate of sodium, or, instead of the calomel, a pill

¹⁸ Berliner klin. Woch., No. 17, '97.

¹⁹ Indian Med. Rec., July 1, '98.

²⁰ Med. Rec., Feb. 6, '97.

containing colchicum, colocynth, and calomel; and the taking of medicated baths. Regular bathing is regarded as almost the most important item, and next to it regular exercise, out-of-doors if possible. All forms of meat are allowed at least once a day, or twice if desired, but no vegetable which grows underground and a restricted amount of sweets. In the treatment of chronic gout, in the intervals between acute attacks, the use of mineral baths, with massage and exercise, are regarded as of the greatest value. If there be some local subacute affection present, the exhibition of calomel, followed on the day after by iodide of potassium and colchicum-wine, may be given, and continued for several weeks. An ointment containing iodine and iodide of potassium applied to the joints is often of service, or, if there is much pain, ichthyol ointment from 30 to 50 per cent. in strength may be used instead. In acute gout treatment should commence with a full dose of calomel, followed by doses of 15 drops of colchicum-wine, which may be increased or diminished according to the absence or presence of toxic symptoms. If the pain is very severe, opium in the form of Dover's powder is indicated.

Armstrong²¹ relates that during the last few years he has given trial in various cases of gouty arthritis and recurrent renal calculi to a method of treatment based on the prolonged administration of only red meat and hot water. Very marked improvement has resulted, which persists in spite of gradual return to an ordinary dietary. Observations on auto-intoxication in relation to the causation of gout and rheumatoid arthritis had led the writer to believe that it is the complex chemical changes brought about by the admixture of red meats with carbohydrates and sugar that causes the ex-

cessive formation of uric acid. His plan is to give the patient a daily allowance of from one to four pounds of lean beef,—steak minced and cooked in various ways,—the patient drinking from one to five pints of hot water, and avoiding all starchy, saccharine, and fermentative articles of food. This treatment is indicated in obstinate chronic gouty arthritis, in recurrent uric-acid calculi, in frequent and intractable migraine, and in cases of persistent gouty dyspepsia. It has proved especially useful in the presence of symptoms of amylaceous and intestinal dyspepsia and of excessive formation of hydrogen sulphide, urates, indican, skatol, creasol, and other toxins. The carbohydrates, by their affinity for oxygen, interfere with the due oxidation of the tissues. Whatever poisonous matters remain in the system are readily eliminated by the taking of the hot water, which also flushes the stomach, liver, and kidneys. The treatment is irksome and trying and must of necessity be carried out with great strictness; therefore its use should be confined to the more difficult cases. It should be prescribed but rarely, and then only under the most careful supervision in cases in which the heart or kidneys are diseased. Used with due care, it is a most efficient and brilliant addition to the therapeutic measures.

In the opinion of H. C. Wood,²² there are three great manifestations of the same thing, universally allied. They are rheumatoid arthritis; podagra, or true gout; and articular rheumatism. One must not attempt to treat gout, but treat the person who comes before him. There is diet for the gouty, but there is a diet for the person. Nevertheless, in

²¹ Brit. Med. Jour., May 1, '97.

²² Med. Rec., July 10, '97.

the large majority of cases sugars and starches must be cut off. But in spare gouty subjects farinaceous diet may be essential. Milk probably suits the largest number of gouty patients. Patients who can take but little exercise at first can gradually be led up to the point of taking a great deal of exercise, and this is essential for further attacks. Strontium salicylate is less disturbing than salicylate of sodium. In some instances it agrees better with the patient when combined with digitalis and strychnine. Medicines, however, will not eradicate the diathesis.

George W. Tobias²³ says that an elegant and perfectly-safe preparation of the drugs, and one which in his hands has never failed, is colchi-sal (colchicine-methyl salicylate). This drug is dispensed in capsules of 20 centigrammes and each contains a quarter of a milligramme of the active principle of colchicum.

F. Levison,²⁴ believing that the uric-acid output does not depend upon the intake of albuminoids, but rather that the disease is secondary to renal degeneration, which, in turn, is caused by the failure of the uric acid to remain in solution, and further noting that the nucleins increase the production of uric acid, would exclude from the dietary all substances rich in nuclein, as thymus gland, liver, kidney, and pancreas. Alcohol, especially in high-percentage liquors and in concentrated solutions, not only increases uric acid, but also directly injures the kidneys, and therefore should be forbidden. Coffee seems to increase the amount of uric acid. In order to prevent the precipitation of uric acid in the kidneys, large quantities of fluid should be taken; the best is boiled water or milk. Mineral acids are not allowed, but vegetable acids, especially

those which are oxidized to alkaline carbonates, can be used without harm. Excessive alkalinization of the urine leads to two dangers: (1) the formation of a phosphatic calculus, or (2) in patients whose kidneys are already diseased and in whose blood there is already commenced a storing up of uric acid, there supervenes an acute attack of gout from an excessive amount of sodium salts in the blood and tissues. So long as the reaction of the morning urine is only feebly acid and the sediment obtained by centrifuge contains no uric-acid crystals, all is attained that is attainable from the use of alkalies, and further increase of the dose is not only unnecessary, but may be directly harmful. Recognizing the dangers of excessive use of the alkalies, von Noorden makes use of calcium carbonate in the treatment of gravel. Calcium carbonate is not eliminated by the kidney, but in the intestines unites with phosphoric acid and is eliminated as calcium phosphate. Thus phosphoric acid being diminished in the urine, the triple phosphates are then more largely formed, while the monosodium phosphate is in a smaller amount; so that the urine remains acid and yet holds uric acid in solution. For acute gouty attacks the greatest reliance is placed upon colchicum. For the stiffness and deformities various methods have been in use—dietetic regulations, massage, iodine-painting, and especially various warm baths.

Recently Grawitz makes use of hot sand-baths; the dry sand is heated to a temperature of 104° to 122° F. and the limb wrapped in it. This has not, according to the author, benefited any instance of arthritis, although it relieves

²³ Kansas City Med. Rec., Mar., '97.

²⁴ St. Petersburg med. Woch., Nos. 1, S. 1; S. 9, '97.

nerve-pains, as in neuritis after influenza.

Of much greater value is the electrical treatment, making use of a battery of forty-eight large Leclanché cells for thirty minutes. The positive pole, connected with a carbon electrode, is immersed in a 2-per-cent. solution of lithium chloride which has been made alkaline with lithium carbonate. The negative carbon electrode is placed in a very weak sodium-chloride solution. The portion of the body to be treated is placed in the lithium-bath and a convenient part, as the hand or foot, is put into the salt solution. A current of 20 to 30 milliampères is now passed, although some patients cannot bear more than 10. Of the 15 patients who received this treatment, 10 were markedly benefited, while in the remainder the results were less favorable or negative. The results claimed are that stiff and useless joints become movable and serviceable; that the pains are completely removed, and the muscular atrophy disappears. The theory of the action is that lithium, at the positive pole, is separated from the lithium-chloride solution and carried into the body in an available form; so that readily-soluble lithium urate is formed from the deposited urates.

R. Newman²⁵ concludes that there is a variety of causes and symptoms of gout. The diet, and treatment, etc., cannot be stated as a routine for all cases.

Static electricity is the best treatment in hereditary gout, and will prevent attacks, if used judiciously at the right time. Static electricity and other electric currents will cure many of the other varieties of gout. Static electricity is generally diffused in the body, and penetrates deeply through tissues and joints.

It acts as a general tonic. The breeze allays any pain, in most instances, in five minutes. In very painful affections of the joints it needs several applications before the pain and infiltration are removed; but, when an attack is in progress, after three applications in a single day freedom of motion and cessation of pain should be expected. It replaces exercise and acts as passive motion.

STERILITY.

Etiology.—G. Pujol²⁶ has investigated fully the question of uterine fibroids and sterility, and finds that all agree as to the frequent co-existence of uterine fibroids and sterility; but all are not at one as to the kind of relationship, the majority regarding the sterility as the result of the tumors, some looking upon the fibroids as the consequence of the sterility. The writer believes that a sufficient explanation of the sterility is found in the various modifications which the fibromyomatous neoplasms produce in the uterus that contains them. These are chiefly metritis (especially granular), inflammatory affections of the tubes, and atrophy in the ovaries of the Graafian follicles; less efficient are the uterine hæmorrhages or hydrorrhœa, spasmodic contractions of the uterus, its displacements, and slight attacks of pelvic peritonitis and the occasional existence of vaginismus as a complication.

In an article on sterility²⁷ E. Rode speaks of a minor abnormality of formation which consists in a prolongation of the posterior column of the vagina on to the hymen, whereby the hymen is

²⁵ Med. Rec., Dec. 11, '97.

²⁶ Arch. de Gynec. et de Tocol., Sept. to Dec., '96.

²⁷ Tidskrift for den norske Lægeforening, '96.

rendered very resistant. A simple remedy is that of a transverse incision of the columna behind the hymen, followed by a suture so contrived as to convert the wound into an anterior-posterior one.

Vedeler²⁸ reports his investigation of 310 sterile women, all of whom had been married more than one year. Seventy-two had been married ten or more years, and the average duration of married life in the remaining was three years. The examination of 50 of the husbands of these women determined that 38 surely had had gonorrhœa, and it was also determined that 34 (68 per cent.) of these had infected their wives. He says it can be concluded that the result of the examination of these 50 sterile marriages be correct, then 235 of the 310 husbands must have had gonorrhœa, and also that 210 wives were infected. That this last approaches the truth is shown in that in 198 (44 per cent.) of the 310 women examined the same inflammatory signs of gonorrhœa were found as in the 34 whose husbands undoubtedly had had gonorrhœa.

Benzler²⁹ made careful and reliable inquiries from 31 men, who were at a previous time of their military service afflicted with epididymitis bilateralis, and who later on married. Of these 38.7 per cent. proved to be sterile, 61.3 per cent. produced posterity—some of them quite a numerous one. In the latter series of men the epididymitis was only in 3 instances associated with vasitis, while in the cases which led to sterility this complication existed in 50 per cent. of the cases. The prognosis in cases of bilateral epididymitis regarding the potentia generandi is, therefore, much more favorable than it is commonly believed to be.

Treatment.—Jones, of Edinburgh,³⁰ states that belladonna is followed by

more or less benefit in every disease to which the female sexual organs are liable; and in married women who, though apparently enjoying the best of health and never suffering from any irregularity of the sexual organs, are yet sterile, the exhibition of belladonna internally for some weeks is so frequently followed by pregnancy as to preclude considering the occurrence as a mere coincidence. During the exhibition of the drug, the external genitals become more relaxed, and the os and the cervix more pliable and softened.

W. Gill Wylie³¹ describes the method he employs in the treatment of dysmenorrhœa and sterility as follows: "When a patient comes to me suffering from dysmenorrhœa and sterility, if the os is sensitive (and this is an indication for the tube), after excluding disease of the Fallopian tubes and other complications, I put her under ether, curette the uterus thoroughly after dilating freely, but not enough to split the cervix, and introduce the fenestrated drainage-tube. I also insert a retroversion pessary to keep the tube in place by holding the os uteri backward. No gauze is employed. A vulvar pad is applied and the patient put to bed, where she is kept quiet for a week. She is then allowed to get up and move about her room for several days. The tube is then removed, and, if she menstruates without pain at her next period, nothing more is done. If she does have pain, several months later I curette again and introduce a smaller tube, keep the patient in bed for a week, and then let her go home wearing the tube, enjoining her to keep the parts

²⁸ Centralb. f. Gynäk., No. 26, '97.

²⁹ Deutsche milit. Zeitschr., No. 4, '97.

³⁰ Columb. Med. Jour.; Mass. Med. Jour., Aug., '98.

³¹ Amer. Gyn. and Obst. Jour., June, '98.

clean by an occasional douche. After the patient has passed through two or three periods, the tube is removed. Of course, all cases in which there is tubal disease are excluded. I have never had a case in which salpingitis or sepsis followed this treatment, although I have been employing it for ten years. I can take ten cases of typical dysmenorrhœa, with anteflexed uteri, and in a few weeks I will cure half of them; in a few months nine out of ten can be cured. In rare cases the treatment has to be repeated a third time. In sterility the results of this treatment have been marvelous. Women as old as 39 have been cured of sterility. The treatment is safe and successful."

YELLOW FEVER.

Diagnosis.—John Guitéras³² states that there is no acute febrile disease in which there are so many signs that may be called pathognomonic. The diagnosis of the disease rests upon three such symptoms, namely: the facies, the albuminuria, and the want of correlation between the pulse and temperature. The facies are extremely characteristic. The appearance of the face is like that of measles before the eruption breaks out, with a more or less pronounced icteroid hue. In the first twenty-four hours, or forty-eight, it is by no means a distinct jaundice. The icteroid hue is often better seen at some distance from the patient than when the eye is closely inspected. In severe cases and on the second and third days of the disease the jaundice becomes more prominent, and later on it may be well marked. The mind is usually clear and there is a peculiar alertness and watchfulness that is not seen in other acute febrile diseases. The albumin appears in the urine usually on the third or fourth day

of the disease. In many cases it is only a trace, but even then by a careful centrifugation granular casts may be found in the urine. The want of correlation between the pulse and the temperature may be a rather late manifestation and may be absent, especially in children. The characteristic feature is that quite often we find that, at the same time that the temperature may be rising, the pulse will be falling.

S. E. Archinard and R. S. Woodson³³ have discovered an agglutination-test to be used in the diagnosis of yellow fever. A drop of blood is taken from the lobe of the ear of the patient and dissolved in twenty times its volume of sterilized water. This is then placed in a culture-tube containing yellow-fever germs which have been active and increasing for twenty-four hours. In from five to thirty minutes after the drop of suspected blood, dissolved in twenty times its volume, is put into the culture-tube the germs in the blood become agglutinated and motility ceases entirely, which shows that the blood is that of a yellow-fever patient. If, however, agglutination does not take place, yellow fever is not present. The yellow-fever germ has been found in between 87 and 90 per cent. of the cases in which this test has been employed.

In a valuable monograph entitled "Yellow Fever: its Nature, Diagnosis, Treatment, and Prophylaxis," etc., by Officers of the United States Marine-Hospital Service, '98, R. D. Murray states that, differentially considered, dengue has a demonstrable rash in the fauces always, between the shoulder-blades generally, and often over the big joints and on the trunk. The pains of

³² N. C. Med. Jour., Dec. 5, '97.

³³ Med. News, Feb. 5, '98.

dengue are in the bones and joints. A dengue patient is in pain and cannot lie still—he does not want to get up. Yellow-fever pains, except the head, are in the muscles, and the patient after four or five days is comfortable in bed, but wants to get up and work.

In malaria the symptoms usually appear after some days of malaise, loss of appetite, discontent, and a general tired feeling. Malarial fever nearly always appears in the day-time or when the victim is at work, and is ushered in with a positive chill. Constipation is the rule, but not so marked a feature as in yellow fever. The malarial tongue is swelled, tooth-marked, and heavy coated, with white edge and yellow or dirty top-area. A yellow-fever tongue is rarely indented; the tongue of the former soon shrinks and has a red edge and red tip, the red tip being diamond-shaped. Herpes does not occur in yellow-fever cases; it is common in malaria. This is, however, a late sign.

Etiology.—Eugene Wasdin³⁴ believes yellow fever essentially an air-borne infection, the entrance of which into the system has been supposed to be by way of the alimentary canal, the upper intestine serving the purposes of incubation of the causative germ, the absorption of its poison giving rise to the disease.

More recently it has been advanced that probably the germ of yellow fever enters the general circulation through the respiratory organs in some obscure manner, and, incubating in the blood, directly poisons this life-giving stream.

According to H. R. Carter,³⁵ all places within a yellow-fever infected district, or town even, are not infected or are infected in unequal degrees. The infection is especially confined to the habitations of men and their environment,

and is conveyed a short distance, possibly 220 metres down the wind, from an infected focus. Two hundred and twenty metres is the maximum distance this infection can be conveyed. This observation (Melier's) is altogether exceptional, and less than half that distance covers the distance to which the infection is conveyed from a single focus.

The infection is heavy and hangs and spreads near the ground. It is unable to pass a close wall of any considerable height, although under the shady side of such a wall it may spread well when once started. It seems especially active at night, and certainly, out-of-doors, is less apt to be contracted on clear dry days.

The rate of propagation of out-door infection is increased in cities in dusty weather. Strong, steady winds in clear weather lessen the infection. There is no reason to believe that yellow fever as usually propagated in this country is water-borne—the fresh-water tanks of infected vessels have never been and are not now emptied at our maritime quarantines.

Edwin Klebs³⁶ makes the remark that yellow fever is transported by sick people, not by goods and not by water; the personal contagion is not direct or confined only to a very near contact; the contagion, emanating from a person, must be deposited in his surroundings before it can infect other persons. Disinfection of the sick and their surroundings is fully sufficient to destroy the germs and check the disease.

Bacteriology.—Havelburg,³⁷ having found in necropsies of yellow-fever pa-

³⁴ "Yellow Fever," etc., by Officers of U. S. Marine-Hosp. Service, '98.

³⁵ *Ibid.*

³⁶ Jour. Amer. Med. Assoc., Apr. 16, '98.

³⁷ Berliner klin. Woch., Nos. 23-26, '97.

tients great masses of a special micro-organism in the contents of the stomach, endeavored to isolate it in order to ascertain whether it was pathogenic or not. The alleged yellow-fever bacillus is very small, straight, and is, for the most part, isolated, the cohesion of two organisms being rather rare. The outlines of its extremities are especially well marked. It is easily stained by basic aniline colors, but not by Gram's method. It can be grown on a gelatin plate, forming in twenty-four hours a white spot which in from twenty-four to forty-eight hours becomes larger and assumes the form of a pin's head. The gelatin is not liquefied.

According to Sanarelli,³⁸ there does not exist any lesion truly pathognomonic of yellow fever, although the changes of yellow fever in their entirety constitute an anatomical criterion more clear and better defined than that of the majority of infectious diseases. The cadavers of the victims of yellow fever are either sterile or they are found to be invaded throughout by a mixture of micro-organisms. The specific microbe, to which the name of "bacillus icteroides" has been given, has never been found alone in the autopsies made. It must be sought for in the blood and in the tissues, and not in the gastro-intestinal tube, in which it has never been encountered. In yellow fever, as in typhoid fever, there takes place in the digestive tube an extraordinary multiplication of the coli bacillus, which is found there in a state of almost absolute purity. Upon the result of investigations it may be said the isolation of the specific microbe of yellow fever is possible in only 58 per cent. of the cases. The reasons for this are easy to understand. In the beginning of the disease the "bacillus icteroides" multiplies very

little in the human organism, a very small quantity of its toxin being sufficient to provoke in man the worst type of the disease. In the second place, the toxin, whether by itself or indirectly through the profound lesions it causes, facilitates in an extraordinary manner every sort of secondary infection. The poison of the "bacillus icteroides" instead of being absorbed through the intestinal walls, is fabricated in the interior of the organs and in the blood. This bacillus is a little rod, with rounded extremities, united at best by pairs in cultures and in groups in the tissues, from two to four micromillimetres in length, and generally two or three times longer than it is broad. The best way to demonstrate, not only its presence, but also its special tendency to arrange itself in small groups, preferably in the blood-capillaries, consists in placing in the incubator, at 37° C. for twelve hours, a fragment of the liver taken from a fresh cadaver in order to favor the multiplication of the specific microbe. The yellow-fever bacillus grows sufficiently well in all the ordinary culture-media. In common gelatin it forms rounded colonies, transparent and granular. The granulation of the colony becomes more pronounced, appearing ordinarily as a nucleus, central or peripheral, completely opaque; in time the whole colony grows entirely opaque. It never liquefies gelatin.

The microbe of yellow fever is pathogenic for the greater number of the domestic animals.

Yellow fever progresses in cycles; at first the specific microbe is very scarce in the organs, and it is only at the end of the disease-cycle, whose duration may be established as between seven or eight

³⁸ Med. Rec., July 24, '97.

days, that the microbe multiplies resolutely and suddenly invades the entire organism, accompanied almost always by other microbes, probably of intestinal origin.

The "bacillus icteroides," once in the organism, not only determines a general intoxication, but also produces specific alterations, which have their seat of election, above all, in the kidneys, the digestive tube, and the liver.

The "black vomit" is due to the action of the gastric acid upon the extravasated blood in the stomach. The vomiting itself is directly provoked by the specific emetic action of the toxins of the "bacillus icteroides" circulating in the blood.

All the symptomatic phenomena, all the functional alterations, all the anatomical lesions of yellow fever are but the result of the action, eminently steatogenic, emetic, and hæmolytic, of the substance manufactured by the "bacillus icteroides."

The diffusion of the virus of yellow fever can take place as well by air as by water.

The "bacillus icteroides," whether by the effect of its specific poison or whether through the grave hepatic lesions which are its most immediate consequence, favors at a given moment the entrance into the organism of septic microbes, which not only end the disease much before the specific agent could do it, but are also prejudicial to the latter, invading at once its domains, suppressing its vegetative faculty and even its vitality.

It is on account of this that these phenomena of microbic antagonism between the yellow-fever bacillus and the micro-organisms of septic infections, instead of being useful to the patient, tend to hasten his death.

The probable cause of the mysterious longevity and resistance of the "bacillus icteroides" on board ships is that the common molds of the atmosphere constitute the great protectors of the "bacillus icteroides." In the holds of ships the moist heat and insufficient ventilation should be regarded as indispensable conditions for the growth of the molds, and therefore as indirectly favorable to the vitality of the "bacillus icteroides."

Surgeon-General Sternberg³⁹ says that it is evident from observations that the micro-organism described by Sanarelli is identical with the bacillus X which has been described by the author. Cultures containing the bacillus X produce vomiting, fatty degeneration, and hæmorrhagic enteritis, proving that its action is identical with that of Sanarelli.

E. Klebs⁴⁰ has conducted researches into the anatomical condition of the liver in yellow fever by means of a new method of staining. The sections, which must not be thicker than $\frac{1}{100}$ millimetre, are stained with a solution consisting of 7 parts of the author's parafuchsin-kresol solution mixed with 3 parts of a concentrated solution of methylene-blue in 5 per cent. borax solution and 3 parts of 1-per-cent. methylene-green. Decoloration is effected carefully by alcohol or by aniline-oil and xylol. The blue color of the section must not be eliminated. In specimens so stained the author finds, between the lines of the bluish tinged liver-cells, intensely-stained red masses, oftentimes forming stripes and masses larger than the liver-cells. The last are transformed and nearly destroyed by fatty degeneration in the median parts of the acini, and compressed by the red

³⁹ Med. and Surg. Reporter, Nov. 6, '97.

⁴⁰ Jour. Amer. Med. Assoc., Apr. 16, '98.

masses, whereas in the centre and peripheral parts only isolated red spots are disposed between the liver-cells. The red masses consist of round, oval-shaped, or irregular balls which will not conglutinate.

With higher powers, one remarks two constituents of these red masses, very deep stained, round, oval, or egg-shaped bodies, and slightly-stained masses, surrounding the first. Oftentimes in one red mass are included two or more of these bodies, somewhat larger than a human red blood-corpuscle, but oval-shaped and of quite homogeneous structure. It is not difficult to demonstrate that the greater, slightly-reddened masses are no other than very enlarged and, in their coloring qualities, deeply-changed leucocytes. Oftentimes they contain the blue-stained nucleus, somewhat altered, elongated, or otherwise deformed. Not all of these red bodies are included in cells, but are found free between the liver-cells, and here the largest forms, which measures more than 13 micromillimetres in length and 12 in breadth, mostly egg-shaped, with one broader and one smaller pole, are found. These contain, mostly, a greater or smaller number of vacuoles and brownish pigment. These larger bodies are not alone situated between, but also in the interior of the liver-cells. The sharply-defined form of these bodies, their different staining, their disposition in the interstitial tissue and in the liver-cells, their forming of vacuoles and pigment in the more advanced stages, certainly indicate them to be parasites of the class of protozoa. In the stomach and duodenum the same oval-shaped and red-staining bodies were always present. Certain blackberry-like forms, circular groups of small round bodies, not present in the liver, are found there also, and

it is suggested that they represent sporulation. It is likely that the source of the disease is, first, a true gastro-duodenitis, remaining such in the milder, lingering, or endemic cases of some countries, but becoming epidemic with the migration of protozoa into the liver.

Pathology.—In the report of the Cuban Commission of Mississippi⁴¹ the diagnostic findings of yellow fever post-mortem were as follow: The skin and sclerotics markedly icteric; usually early ecchymotic spots on the back of the neck, shoulders, and lumbar regions, thighs, calves of legs, and the ears. The abdomen is usually dry, sometimes a little fluid is present; the liver is contracted away from the ribs, of a box-wood color, bloodless, and friable. The gall-bladder is contracted, usually empty, but sometimes containing a thick, tarry fluid. The spleen is normal in size and color. The kidneys are normal, sometimes showing signs of recent acute inflammation. The stomach is usually anæmic, generally shows exfoliated spots where hæmorrhages have occurred, and frequently contains black vomit. The intestines generally contain a pasty-colored material like the stools, only not so black. The mucous membrane of the whole alimentary tract shows the most decided effects of the poison. The heart often shows traces of fatty degeneration.

According to R. D. Murray,⁴² the primary lesion of yellow fever is in the duodenum and the symptoms of the disease follow generally in regular order. If the stools could be all and carefully examined some time a mass of white

⁴¹ Jour. Mississippi State Med. Assoc., May, '98.

⁴² "Yellow Fever," etc., by Officers U. S. Marine-Hosp. Service, '98

mucus with a black or brownish centre will be found.

Prognosis.—Geddings⁴³ states that probably no case of yellow fever can occur without presenting albuminuria at some time, though that time be limited to a few hours, perhaps. The quantity present on first detection, and its increase or diminution from day to day, form, perhaps, a fairly good guide to prognosis. If it appears, increases gradually, and then begins to diminish, prognosis is good. If, on the contrary, it appears at first in large amount, persists or increases abruptly, trouble may be anticipated. The two gravest symptoms that can arise during the course of a case of yellow fever are undoubtedly black vomit and suppression of urine. Partial or complete suppression of urine is of the gravest accident that can happen in the course of yellow fever.

Treatment.—Sanarelli⁴⁴ says that good results from serum-antiamari are difficult to obtain if the treatment be applied when the disease is already advanced, when the "amarillic" poison largely accumulated in the organism has already induced those grave anatomical and functional changes which the serum cannot undo and which of themselves suffice to cause death. We must for the present, therefore, restrict our application of the serum exclusively to the first period of the malady. It can be injected into patients in the desired quantity up to the moment when all hope of success is not yet definitely abandoned. One should begin with a dose of 20 cubic centimetres, and, if appreciable improvement fail to set in, a second, a third, and even further doses may be injected, always being guided by the patient's resisting power, or his general condition, or the "period" of the malady, or the complications. The injections

must be practiced subcutaneously in the region of the thighs or nates, but in urgent cases it is preferable to introduce the serum directly into the veins. These injections must be practiced under antiseptic safeguards of the most stringent kind.

E. K. Sprague⁴⁵ remarks that the apparent lack of success that has thus far attended the treatment of yellow fever with antiamarillic serum constitutes no argument against the bacillus as the cause of the disease, as there are yet many diseases in which the microbic cause is incontestably established, but for which we are still unable to procure a specific curative serum.

H. M. Folkes⁴⁶ says that a case of yellow fever can be treated in a room in a crowded hotel, and by taking the following sanitary precautions no other guest in the place need become infected: Over the windows should be placed a double thickness of mosquito-netting or some such material, kept constantly moist with a 1-to-500 bichloride solution. The patient's gown and bed-clothing should be removed twice daily, immediately putting the same in a 1-to-500 bichloride solution, and a rubber sheet placed next to the mattress. The fewest possible things should be allowed in the room, and these are to be wiped with bichloride solution once daily. All dejecta, sputa, etc., from the patient are to be placed in the same solution, and all cups, towels, glasses, etc., are to be treated in like manner. The nurse should stay in the sick-room, or else,

⁴³ Annual Report Marine-Hosp. Service, '94; "Yellow Fever," etc., by Officers U. S. Marine-Hosp. Service, '98.

⁴⁴ Lancet, Mar. 26, '98.

⁴⁵ "Yellow Fever," etc., by Officers U. S. Marine-Hosp. Service, '98.

⁴⁶ Ga. Jour. Med. and Surg., July, '98.

when going out, she should take a bichloride bath, if possible, and put on sterilized clothing all over,—her shoes included,—unless she is simply leaving for a few minutes, in which case she should take the same precautions as the physician.

The physician should always put on a cotton gown wet with the same solution before going in the sick-room; this is to be removed when he comes out, and his hands and face should receive a formaldehyde or bichloride bath at once.

On recovery or death of the patient, if the room and its contents are thoroughly disinfected it will positively prevent a spread of the fever.

H. D. Geddings⁴⁷ believes that the treatment of yellow fever should be symptomatic and directed toward meeting plain and specific indications. For the initial stage, as well as for properly initiating a systematic treatment, administering a hot foot-bath containing mustard is excellent. As soon as the patient is made comfortable in bed, a sharp purge, preferably mercurial, should be given; calomel, 5 grains; compound powder jalap, 10 grains; administered in capsules found most efficient. Should this fail to move the bowels freely within six or eight hours it may be followed by a moderate dose of castor-oil, a Seidlitz powder, or a bottle of citrate of magnesia.

Closely following the first purgative should be administered one of the coal-tar febrifuges; phenacetin, $7\frac{1}{2}$ grains, or antipyrin or antifebrin, 10 grains, either of which may with advantage be combined with $1\frac{1}{2}$ to $2\frac{1}{2}$ grains of citrate of caffeine. Repeated doses of the antipyretic are not needed, nor, indeed, indicated. One, two, or, at most, three doses in the first twenty-four hours of the disease will accomplish all that

is to be gained from this series of remedies.

The gastric irritability may be controlled by sinapisms to the epigastrium, abstention from fluids, and frequent ingestion of small pieces of ice. Should nausea or vomiting persist, the administration of cocaine hydrochlorate, in doses of $\frac{1}{4}$ to $\frac{1}{2}$ grain every hour or two, will often act almost magically. Small quantities of carbonated beverages, as Vichy, seltzer or apollinaris water, ginger-ale, or very dry champagne, administered ice cold, will often prove of service. Considerable relief is also derived from the application to the epigastrium of a liniment composed of olive-oil and menthol.

R. D. Murray⁴⁸ gives three or four compound cathartic pills at once and as soon as possible, a hot foot-bath, with or without mustard and salt. As soon as possible, if fever is above 102° , any coal-tar derivatives in $7\frac{1}{2}$ -grain doses, with some bicarbonate of soda and caffeine to be given. After the bath and a good sweating, under blankets, for from four to six hours, the patient should be rubbed dry and covered with two blankets. The coal-tar derivative should be repeated every three to six hours if fever keeps above 102° . Orange-leaf tea, apollinaris water, lemon-grass tea, hot lemonade, ginger-ale, small sips of ice water, and other drinks may be given *ad libitum*. No spirits of any combination should be given. If the bowels are not freely and comfortably relieved within six hours, a small saline is administered. After thirty-six hours an enema should be given every day.

⁴⁷ Annual Report Marine-Hosp. Service, '94; "Yellow Fever," etc., by Officers U. S. Marine-Hosp. Service, '98.

⁴⁸ "Yellow Fever," etc., by Officers U. S. Marine-Hosp. Service, '98.

H. D. Geddings⁴⁹ states that when "black vomit" appears the treatment should be directed toward the general hæmorrhage diathesis. The most efficient remedy is found in the tincture of the perchloride of iron, 15 to 30 grains every hour or two, or, if the vomiting is frequent, after each act of emesis. Counter-irritation to the epigastrium, the administration of stimulants, preferably champagne or good brandy, administered in carbonated water and given cold, swallowing of ice, and administration of cocaine make up about the sum of remedial agents.

In suppression of urine, counter-irritation over the region of the kidneys with turpentine or mustard, dry cups, the application of hot-water bags, all should be tried. A *tisane* of watermelon-seed is efficacious alone or given in combination with spirit of nitrous ether.

Frequent washing out of the lower bowel with enemata of warm water and soap is very important. A well-oiled rectal tube should be passed as far up into the bowel as possible, and with a fountain-syringe elevated not more than a foot or two 2 or 3 pints of warm, soapy water should be slowly forced into the bowel. In malarial regions it is a good practice to administer 30 to 45 grains of quinine or cinchonidia in the first twenty-four hours, exhibiting the drug per rectum if the stomach is irritable.

As regards diet, Geddings thinks that the yellow-fever patient should be well nourished, but the most scrupulous care should be exercised in the selection and administration of food. "A little and that often" should be the rule. For the first few days milk with lime-water given cold, then animal broths, concentrated, but free from fat, should be the regimen. The fever being reduced, soft-

boiled eggs, milk-toast, and small bits of the white meat of chicken and tenderest steak may be permitted. Probably at least ten days or two weeks should elapse before the convalescent, by the easiest stages, should be permitted to resume ordinary diet.

Prophylaxis.—H. R. Carter⁵⁰ states that as far as the selection of a living place is concerned, non-infected location, as far from any known focus of infection as possible, or residence portion of the town liable to be or become such a focus, should be preferred. It should, if sufficient distance be unattainable, be to the windward (prevailing wind) of such portion of the town or separated from it by trees, etc., and be located on high, well-drained ground, as much exposed to wind as possible, and not so shady as to be damp. The place chosen for residence should be kept dry by means of ditches. It should be kept very clean, and free from vegetable as well as animal matter, decaying leaves and wood, especially sawdust. The houses should be preferably built of wood with free circulation of air under them, light and wind penetrating everywhere. Articles stowed away and handled in the non-resident portion (business district) of the town are little liable to be exposed to infection. The capability of an article to retain infection depends upon the nature of its exposed surfaces. If these be hard, smooth, and non-absorbent, the article will be little likely to convey infection. Household goods are most exposed to infection and should be burned.

One should not enter houses unneces-

⁴⁹ Annual Report Marine-Hosp. Service, '94; "Yellow Fever," etc., by Officers U. S. Marine-Hosp. Service, '98.

⁵⁰ "Yellow Fever," etc., by Officers U. S. Marine-Hosp. Service, '98.

sarily; should not go out of uninfected quarters at night; should not receive or come in contact with any fabrics or household goods about the exposure of which to infection there is doubt.

There are certain personal factors affecting susceptibility which we cannot influence, and yet of which we may avail ourselves advantageously, as

(a) The lessened susceptibility of the negro race.

(b) The lessened susceptibility of those living for years (better for generations) under the conditions which obtain in the tropics.

Thin, spare, vigorous men are less liable to develop the disease, and bear it well.

Most potent auxiliaries to an infection are,

1. A sudden chilling of the surface, especially if wet with perspiration.

2. Excessive exposure to the direct rays of the sun.

3. Excessive physical fatigue.

4. Anxiety and mental distress generally, especially fear of the disease.

The constipation nearly always associated with the advent of the fever probably has an etiological influence.

A meat diet is claimed to increase the susceptibility to the disease, as well as to make it more severe. The same is true of alcoholics. The free use of water and a diet of juicy fruits and fresh vegetables, should put a man in a better condition to go through the disease and probably lessen his chance of developing it.

To avoid chilling the surface, flannel should be worn.

When the disease has invaded a town, the author argues that if there be but few foci, and these be known, the chances of suppressing the fever are

good. If, in addition, all who have been exposed to infection are known and can be properly provided for, the chances of success in arresting the disease are greatly increased.

If the fever be confined to one section of the town, even if pretty general therein, it may be possible to so isolate that part as to preserve the remainder. The patient should, if possible, be moved to an isolated place or a well-appointed hospital. Removal during the first forty-eight or sixty hours prior to the "stage of calm" is not specially injurious. After that time it is to be deprecated. If the patient is moved, all possible precautions to prevent infection of his new quarters must be taken. Cleanliness, dryness, good ventilation, and sunshine are all important. No fabrics, carpets, hangings, etc., not absolutely necessary should be allowed in the room. The clothing, bedding, etc., which go with him if moved must be immediately disinfected. A rubber sheet to protect the mattress must be placed on the bed. The bed-linen and shirt must be changed daily, oftener if soiled; and the rubber sheet be changed when necessary. All fabrics used about the patient should at once be put into an antiseptic solution. This should be done in the patient's room and the floor should be wiped up daily with a similar solution. All excreta should be disinfected or destroyed. The physician should wear linen or other smooth clothing, or change it if he goes out. These precautions are recommended only when there are very few patients and every real risk, however slight, is to be avoided. Until premises are released from observation they must be under guard. The premises adjacent to those of the patient which, from propinquity, communication, or direction of wind can reason-

ably be judged to have received infection are also to be disinfected.

The inmates of the house of the patient (unless immune to yellow fever) should be removed from the house, all clothing, etc., disinfected and kept under observation—"quarantined," in a place free from infection and so situated that if any of them sickens he may not establish a focus of infection dangerous to the community.

It is to be noted how rarely people taken from infected premises and placed in camps, or under the conditions of camp-life, develop fever.

When the fever cannot be suppressed, the providing of a legitimate means of egress, if safe, is an added safeguard, and an important one, against the infection of clean territory.

Stress should be laid on early depopulation. Classes who may leave are those who have been certainly not exposed to infection. The others may go (1) directly to points incapable of receiving the infection of yellow fever, generally northern points—high altitudes—to remain there indefinitely, or for a time to cover their incubation; (2) to points capable of receiving such infection but through a camp of detention.

There seems no reason why baggage going north should be disinfected. Indeed, there are good sanitary reasons for not doing so. Every obstruction, however slight, put in the way of people leaving an infected town to some extent prevents their leaving and to a disproportionate extent induces them to put off leaving. The sooner they depart the better.

W. F. Brunner⁵¹ recommends that businessmen leave the city every afternoon before sunset and spend the night at some of the small towns near by, returning next morning after sunrise to

pursue their different callings. In a seaport or in a town on a river persons should be prohibited from sleeping near the wharf or river-front.

Care should be taken to sterilize all clothing and material in an infected room and to thoroughly disinfect the room and contents after recovery or death of the patient. One of the principal precautions to be taken is in the matter of clothing. Flannel, light in texture and color, is the best material.

Wasdin⁵² claims that the observance of the rules for personal cleanliness is imperative. All undue fatigue should be avoided, and the question of suitable clothing carefully considered. The greatest regularity in taking meals should be observed and the use of boiled water is advisable. Alcoholic beverages to one unused to them are harmful; to those habituated, their use in moderation seems necessary. The use of internal medication to ward off this disease is useless. The hygiene of environment is the more important. One should select a dry, well-drained abode, to which sunlight has free access, and which can be thoroughly ventilated. One should live such a way that the precautions suggested above may be intelligently exercised, and he will have done all that a sound mind within a sound body can do.

Seaton Norman⁵³ thinks that the advisability of depopulation depends upon the size of the town or village, the density of the population, and the number exposed.

J. H. White⁵⁴ recommends the following method of disinfection:—

⁵¹ "Yellow Fever," etc., by Officers U. S. Marine-Hosp. Service, '98.

⁵² *Ibid.*

⁵³ *Ibid.*

⁵⁴ *Ibid.*

1. Apartments or dwellings infected with yellow fever to be disinfected by one or more of the following methods:—

(a) By a thorough washing of all surfaces of apartments with an efficient germicidal solution.

(b) By sulphur dioxide for twenty-four hours' exposure, 4 pounds of sulphur for each 1000 cubic feet, plus due allowance made for waste.

(c) By formaldehyde-gas in not less than a 4-per-cent. per-volume strength, and not less than six hours' exposure.

[One litre of 40-per-cent. solution of formaldehyde-gas will evolve about 1.425 litres (50.1 cubic feet) of gas at 20° C. (68° F.).]

2. Grounds, out-buildings, etc., deemed to be infected, to be disinfected with a strong solution of crude carbolic acid (carbolic acid, crude, 2 parts; sulphuric acid, 1 part; water, 25 parts) or an acid solution of bichloride of mercury (1 to 500); disinfection of ground, preferably by fire.

3. Bedding, wearing-apparel, carpets, upholstered furniture, and the like to be disinfected by one or more of the following methods:—

(a) By steam at a temperature of 212° to 216° F.; thirty minutes' exposure.

(b) By boiling, all parts of the articles to be surmerged.

(c) By saturation in an efficient germicidal solution.

(d) By thoroughly wetting the surfaces of articles with a 40-per-cent. aqueous solution of formaldehyde, and placing them in a closed space for not less than twelve hours.

(e) Where surface-disinfection is required, formaldehyde-gas of not less than a 4-per-cent. per-volume strength and not less than six hours' exposure, or

by sulphur dioxide for not less than twenty-four hours.

4. The dejecta from cases of yellow fever to be disinfected by an efficient germicidal solution.

Mails to be disinfected by one of the following methods:—

(a) By formaldehyde-gas.

(b) By sulphur dioxide.

(c) By steam.

(Newspapers must be made up in such packages as shall be penetrable to the disinfectant used.)

Articles injured by steam, such as rubber, leather, and containers, to which disinfection by steam is inapplicable, to be disinfected:—

(a) By thoroughly wetting all surfaces with an efficient germicidal solution, the articles being allowed to dry.

(b) By exposure to sulphur dioxide.

(c) By exposure to formaldehyde-gas.

The application of gaseous disinfection to these articles should be made in a closed space, air-tight, or as nearly so as possible.

The following are considered efficient germicides:—

1. Bichloride-of-mercury acid, 1 to 1000.

2. Carbolic acid, pure, 5-per-cent. solution.

3. Trikresol, 2-per-cent. solution.

4. Solution of formaldehyde, 1 to 500 (which is 1 part of a 40-per-cent. solution of formaldehyde to 199 parts of water).

5. Solutions of hypochlorite of calcium (chloride of lime).

Post-epidemic Disinfection.—In the work of post-epidemic disinfection, Surgeon-General Walter Wyman⁵⁵ instructs

⁵⁵ "Yellow Fever," etc., by Officers U. S. Marine-Hosp. Service, '98.

his officers to carry out the following measures:—

(a) The medical officer in command shall make, or cause to be made, house-to-house inspection of all infected localities, and obtain complete lists (giving number and street when practicable) of all buildings in which yellow fever occurred or where suspicious disease existed during the past summer and fall.

(b) This inspection should be made by competent sanitary officers, under direction of the medical officer in command, and every part of the premises must be carefully inspected.

(c) The inspection includes the inspection of all streets, alleys, and by-ways.

(d) The inspection should include an examination into the water-supply.

A complete list to be made of all persons exposed to, or who may have contracted, the disease, with the result in each case.

It is recommended that after the inspection above provided for has been made the medical officer shall designate a competent sanitary officer to perform the thorough disinfection and cleansing of all houses and premises which he may have decided require the same, said disinfection to begin as soon as practicable

after the inspection referred to has been made in any locality.

It is recommended that the removal of all refuse, garbage, and other deleterious matter be included in the work of disinfection.

It is recommended that the inspectors inform all parties whose houses are visited that no injury to their houses or contents will result from the disinfection contemplated, and that it is absolutely necessary for the protection of themselves and the community in which they live.

The use of formaldehyde-generators or lamps is recommended for the disinfection of houses and their contents.

Both before and after disinfection the houses should be opened and thoroughly aired—"chilled," if the weather is favorable—and later on all the rooms, closets, etc., should be exposed to several hours' airing during freezing weather, and repeated at intervals during the winter.

Stables, pens, etc., can be disinfected by the use of bichloride-of-mercury solution, 1 to 500, or carbolic-acid solution, 50 parts to 1000 parts (applied by means of a spray), if all exposed surfaces are completely saturated. Privies may be disinfected by chloride of lime or strong solution of carbolic acid.

Cyclopædia of Current Literature.

ABDOMINAL SECTION.

Intestinal paralysis or obstruction is more often the cause of fatal sepsis, either wholly or in part, than *vice versa*, in many cases the septic matter finding its way through the stretched intestinal walls. Exposure of the peritoneum, handling of the viscera, production of raw surfaces, and leaving dead matter

(bloody oozing and *débris*) are followed by intestinal adhesions in from 12 to 36 hours, and these adhesions produce more or less intestinal paralysis and sometimes obstruction. On the day before a peritoneal section, the patient should be purged sufficiently to reduce the gaseous distension of the intestinal coils (that they may be kept out of the way during

the operation), obtaining as many as six or eight large stools, while patients of relaxed fibre should receive full doses of strychnine from the time they come under observation. Two hours before the operation 2 teaspoonfuls of the fluid extract of cascara are given. Immediately on awaking from the anæsthetic the patient receives a drachm of magnesium sulphate every hour; at the end of six hours a stimulating enema is administered and repeated till gas passes between enemas; then the saline is discontinued. In simple operations where undue haste is not necessary the salines and enemas are given a little later. The author presents as presumptive proof of the value of this method a record of 105 consecutive recoveries after peritoneal sections since its adoption. H. T. Byford (*Amer. Jour. of Obstet.*, July 1, '98).

ADENOID VEGETATIONS AND DEAF-MUTISM.

A subject of absorbing interest is the relation that apparently exists between deaf-mutism and the presence of adenoids. Thus, one observer found adenoids in 59 per cent. of the boys and girls who were deaf-mutes. It is probable that, if sufficient attention were given to adenoids in early infancy, a material diminution in the number of these unfortunates would result. E. Fayette Smith (*Phila. Med. Jour.*, July 9, '98).

ADENOIDS, POST-NASAL.

Diagnosis.—In nine-tenths of the cases of mouth-breathing children the lesion will be found to be either enlarged faucial tonsils or post-nasal adenoids, or a combination of both. These conditions must be excluded before the case is put down as nasal catarrh. The symptoms in these cases are always pretty

much alike—usually anæmic and ill-nourished, the child breathes partially or wholly through the mouth, catches cold at all times, suffers from impaired hearing, oftentimes earache; is croupy, with more or less bronchial catarrh through the winter. Such general symptoms as these always indicate more than simple nasal catarrh. The diagnosis is very easily made if one looks into the anterior part of the nose. Although there may be accumulated nasal secretions, no deviation of the septum and no hypertrophy sufficient to occlude the nostrils will be found. Enlargement of the middle turbinate is very rarely a factor in children; so at once one is brought back to the vault of the pharynx as the source of trouble. From clinical experience it can be stated that there is more or less enlargement of the third tonsil in all these cases. Great enlargement of the faucial tonsils occurs more rarely than in former years, but moderate enlargement of the post-nasal tissue is just as frequent.

Diagnosis of post-nasal adenoids and faucial tonsils can frequently be made with Munger curette, which is much more easily introduced than the finger, and anybody who is accustomed to its use can quickly determine whether an enlarged tonsil exists. If the curette is introduced into an adult pharynx the surface will feel as smooth as glass, and it is not possible to sink the point of the curette into the tissues. It is difficult even to scarify a normal post-nasal space, because the surface is so hard and slippery. It is quite different in a vault filled with enlarged adenoids. When the point of the curette reaches this tonsillar enlargement, it will be felt to be soft and spongy, the point of the curette is easily pushed into it, and resistance is felt when one attempts to push the

curette downward. This makes the diagnosis certain. Rice (*The Canada Lancet*, Aug., '98).

ALGINATE OF IRON.

Alginic acid is a new organic acid obtained from algæ. It is a nitrogenous body, and as first isolated it forms a light-brown, gelatinous substance. Alginic acid combines with almost all the bases to form compounds, soluble and insoluble. In addition to these compounds, alginic acid combines with many alkaloids, forming soluble films which may have a very extensive and useful application in the field of medicine. All of the alginate salts are little acted on by pepsin, and therefore pass through the stomach almost unaltered—a point of great practical importance in the therapeutic application of some of them.

Alginate of iron is a tasteless, brown, insoluble powder. Compared with the albuminous ferric compounds, the proportion of iron is large, viz.: 10.92 per cent. The alginate of iron is best administered in a fine powder in doses of from 10 to 15 grains, given thrice daily. Unlike the majority of iron preparations, the alginate has no astringent effect; thus the alginate of iron is a compound containing an unusually large percentage of the element in a very assimilable and active form. Perhaps the fact that it is little acted on by gastric digestion accounts for the tolerance which the stomach exhibits toward it. Indeed, in the majority of instances it proves a gastric sedative, and although given in large doses, instead of inducing or increasing, it diminishes, constipation. William Maclellan (*Glasgow Med. Jour.*, July, '98).

ANIMAL EXTRACTS.

A possible danger to health may be incurred in the use of thyroid and thy-

mus glands, that may be sold by butchers under the name of "sweetbreads." The writer has made some investigation of the subject and finds that there is a confusion of terms, intentional or otherwise, in the minds of the sellers of meat. He says: "In talking to a butcher I was surprised to hear him say that there were three sweetbreads in an animal. I asked him where they were situated, and he told me at the root of the neck, in the cavity of the chest, and in the belly of the animal. I questioned him more closely and discovered that his three sweetbreads corresponded respectively to (1) the thyroid gland; (2) the thymus gland; and (3) the pancreas. To make quite sure that no mistake has been made, I watched him, a day or two after, kill and cut up a bullock. The thyroid gland in this animal is situated low down at the root of the neck, over the trachea. The thymus (which he told me was bigger in the calf than in the bullock) is placed in a somewhat similar position to where it is in man; while the pancreas (the most important organ) he regarded as the least significant, and told me it was 'given in with the liver.' He told me, moreover, that if he were asked for calf's sweetbread he would always give (what we call) the thyroid and thymus glands. It seems only right for every physician to be on his guard when ordering 'sweetbreads' for his patients that the pancreas be provided, and not these other glands. An undercooked calf's thyroid gland being repeatedly given in the place of a real 'sweetbread' might produce untold mischief in a patient, as well as perplex the mind of the prescribing practitioner in a peculiar and undeserved manner." Parry (*Canada Lancet*, July, '98).

Whereas some patients can take large doses of thyroid with impunity, others

are injuriously affected by small amounts, and one should begin with a minimum dose,—say 3 grains daily,—and increase it very slowly, watching the heart and kidneys carefully. In cases of bleeding fibroids, thyroid has an influence toward checking the loss of blood, and in certain cases it is followed by diminution in size of the growth. Its use in appropriate doses is followed by improvement in the general health, probably due to the cessation of loss of blood. Nine to 15 grains is the maximum dose, 1 to 5 grains three times a day. The average duration of treatment, nine weeks. W. E. Moseley (*Med. News*, July, 9, '98).

BLENNORRHAGIC ARTHRITIS.

Treatment.—It is advisable in all acute inflammations of the joints to examine the urethra. In 90 per cent. of the cases urethritis will be found. The cases may be divided into four groups: First, where effusion alone occurs; second, where there is formation of fibrin and thickening of the capsule; third, periarticular plegmon with impairment of the action of the tendons and elasticity of the ligaments; fourth, where ankylosis occurs very early. The puncture of the joints and the injection of a solution of carbolic advised. If there is a periarticular affection, the joint should be opened and washed out. König (*Sammlung. klin. Vorträge*, No. 170, '96; *Boston Med. and Surg. Jour.*, July 7, '98).

BOILS, CARBUNCLES, AND FELONS.

Treatment.—The occurrence of suppurative processes should always be regarded as evidence of faulty metabolism, and search should be made to discover and rectify what is wrong. Patients with boils, carbuncles, and felons are

never in perfect health, although it is extremely difficult at times to discover the cause on which the trouble depends. Iron is most commonly needed, but quite as often there will be digestive and assimilative difficulties. In the local treatment of the disease, the objects aimed at are, first, the protection of the inflamed area; second, exclusion of the air; third, a slight antiseptic action. To obtain this end the inflamed surface is covered with a thick layer of absorbent cotton, on the center of which is smeared an ointment of carbolic, ergot, zinc oxide and powdered amyl, made up with an unguent of rose. When pus is present, the skin is left to part spontaneously. The ointment is applied constantly until the carbuncle heals.

In felons, the diachylon, or litharge, ointment, prepared according to the formula of Hebra, is employed; the pain grows less, and the patient's general condition rapidly improves, and the lesion in the finger terminates in a short time in resolution. L. Duncan Bulkley (*Brit. Med. Jour.*, Oct. 2, '97).

BRIGHT'S DISEASE.

Treatment.—There is a good deal of mischief done by iron in Bright's disease. Basham's mixture in Bright's disease was never suggested for any directly curative purpose, but simply as a remedy for the anæmia which is so conspicuous a symptom in many cases, and for this purpose it still is and always will be useful. But not every case of Bright's disease is anæmic, and as iron has no specific curative effect it is clearly not indicated in non-anæmic cases. Nay, more, it is often harmful. It may be laid down as a rule to which there is almost no exception that iron is not indicated, and should not be prescribed in cases of acute Bright's disease. On the

other hand, after the acute symptoms have passed away and convalescence sets in iron is oftēn very useful.

A second class of cases in which iron is contra-indicated is chronic interstitial nephritis, in which it is more promptly and dangerously harmful than in any other form known of Bright's disease.

The form of Bright's disease in which iron is best borne is chronic parenchymatous nephritis. And as this is apt to be associated with more or less anæmia it becomes a most valuable remedy in overcoming this symptom. Even here the doses given are usually needlessly large. The author's practice is to determine the proper dose by an examination of the stools, and if these are decidedly blackened, too much is being given. On the other hand, a slight coloration may be permitted. Basham's mixture is no more diuretic than the bulk of water which constitutes its menstruum. James Tyson (*Jour. Amer. Med. Assoc.*, July 23, '98).

CHLOROSIS.

Treatment.—One of the most promising recent advances in the treatment of obstinate cases of chlorosis is diaphoresis. Better than the dry, hot-air baths for this condition are the hot-water baths. A characteristic symptom in most chlorotics is pain between the shoulder-blades. The erector spinæ muscles suffer from the act of keeping the patient erect. The reflex action of the hot baths causes more blood to be brought to the muscles and relieves this muscular discomfort. A bath at 105° F. with a wet towel wrapped around the head is given for one-half to three-fourths of an hour. This causes plentiful perspiration during the bath. It is followed by a cold douche to prevent

further perspiration, which would be exhausting. A bath is given three times a week for three to four weeks. Rosin (*Med. News*, May 14, '98).

COLOTOMY AND COLOSTOMY.

Colostomy consists in bringing the descending colon up to the anterior abdominal wall, to which it is stitched, the opening into the lumen being made at once or after an interval, according to circumstances.

The great disadvantage of this operation is that it does not entirely prevent the entry of fæces into the distal part of the bowel, where they tend to set up inflammatory troubles. In colotomy, the gut is cut completely across, the proximal portion brought out of the wound, and the distal closed by sutres and returned to the abdomen. This method is not entirely satisfactory, as the distal end tends to become distended by the accumulation of its own secretion, which may eventually lead to ulceration. König and Sonnenburg leave the upper extremity of this portion open and attach the artificial anus to the abdominal wall below, whereby the rectum can, if desired, be irrigated from above. Another means of preventing fæces from getting into the rectum is by the formation of a spur, but the disadvantage of this method and colotomy is that they leave a long and freely movable colon and mesocolon. The best method consists in the ordinary operation of colostomy performed at one sitting, but preceded by partial occlusion of the distal portion of the bowel. A ligature is tied around this, occluding it to about one-half its diameter, and the bulging serous surfaces on either side are sewn together with interrupted stitches. An artificial constriction is thus produced, which prevents the accumulation of fæces in the rectum. In attaching the

gut to the belly-wall, first sew the serous and muscular coats of the intestine to the parietal peritoneum, and then pass the ordinary sutures through both bowel and abdominal wall. If, however, this will lead to considerable tension, attach the intestine to the fascia of the external oblique, leaving the skin free, but shutting off the muscular planes from the risk of infection. Mosetig-Moorhof (*Wiener Med. Presse*, No. 3, '98).

CONDURANGO.

Condurango-bark, which has been extolled for the treatment of various gastric affections, and particularly in gastric cancer, exercises a peculiarly-calmative effect on gastric pains in the dose of $7\frac{1}{2}$ to 12 grains powdered condurango daily, in pill form. It has also been observed that the pains and vomiting of round gastric ulcer, which had resisted cocaine, rapidly ceased on administering $2\frac{1}{4}$ grains, four times a day, in pill form. Lemoine (*Sem. Méd.*, vol. xviii, p. 110).

CONVULSIONS, INFANTILE.

Etiology.—Alcoholism on the part of the nurse is a competent cause of convulsions in a breast-fed child; such convulsions are preceded by nervous irritability, general hyperæsthesia, but without gastro-intestinal derangement, elevation of temperature, or pulmonary complication. They are apt to appear in extremely well-nourished children. As regards the fits, they show marked tendency to increase in number and severity. In some instances there may be anuria. Under such circumstances it is necessary to inquire carefully into the habits of the nurse, and to make a change as early as possible. Meunier (*Jour. de Méd.*, April 25, '98).

DYSENTERY.

Acute tropical dysentery treated with very satisfactory results by the administration of ammonium chloride. The salt was given in drachm-doses every four hours, and the patient placed on milk-and-arrowroot diet. In the majority of cases blood was absent from the stools on the third or fourth day. Ipecac is useless in the treatment of tropical dysentery, and opium should never be given in the early stages; its beneficial effects are only seen in the last stages when combined with cannabis Indica, and when other drugs have failed. J. W. S. Attygalle (*Brit. Med. Jour.*, No. 1949, p. 1197, '98).

ENTEROCLYSMS.

All the necessary apparatus is a rubber rectal tube and a fountain-syringe, or a funnel. A rectal tube of soft rubber, having rather thick walls, and measuring two feet in length by three-eighths inch in diameter, is preferred. The surface of the tube is made very smooth, and the tip is rounded and slightly tapering, and has a terminal opening. The reservoir is usually placed at a height of four or five feet, and the tube is introduced a distance of eight inches. A child is usually given an enteroclysm of 1 pint, an adolescent one of 2 pints, and an adult one of 3 or 4 pints. Ten minutes are usually occupied in administering the enteroclysm, and the fluid should be retained for a like period. The occurrence of colic is an indication of the presence of gas, of the use of too large a quantity of fluid, or of its too rapid introduction. Experiments on the bodies of four children to determine how far up the intestinal tract these injections go showed that there was no difficulty in causing the fluid to pass through the ileo-cæcal valve into the

small intestine, and even out through the mouth and nose.

The following are the conditions in which enteroclysms are chiefly useful: Obstinate and long-standing constipation; autointoxications due to decomposition of the intestinal contents; many forms of irregular gout; cases of chlorosis in which fæcal anæmia is a prominent element; progressive pernicious anæmia occurring without obvious cause; diabetic and uramic conditions; cholera; typhoid fever in the early stages; pseudomembranous colitis, and insolation. Judson Daland (*Phila. Med. Jour.*, July 9, '98).

EXOPHTHALMIC GOITRE.

Treatment.—Good results obtained from the use of the constant galvanic current in the treatment of Basedow's disease. The exophthalmus diminished or disappeared, the general condition improved and there was diminution of the disordered cardiac innervation, and in volume of the hypertrophied thyroid body. Bertran (*Arch. de Ginecol., Obs. y Ped.*, No. 5, '98).

FALLOPIAN TUBES, DISEASES OF.

Diagnosis.—Pyosalpinx is to be suspected if dilatation of the tube follows gonorrhœal infection and if the tumor is very closely adherent. Hydrosalpinx and pyosalpinx are usually double, while hæmatosalpinx is unilateral. If the tubal mass is of large size and there is no extensive adhesion, there is probably a hydrosalpinx; pressure is less painful than in the case of pyosalpinx. While the tumor is still movable it may be mistaken for a small ovarian cyst, and especially for an intraligamentous cyst. The latter, however, is more decidedly lateral, and is not usually separated from the uterus by the space cor-

responding to the pedicle of the tubal cyst. The differential diagnosis from tubal pregnancy in the first four months is almost impossible to make with certainty.

In the majority of cases where these fætal cysts have been removed, the operation has been undertaken for supposed salpingitis. Enlargement of the uterus and expulsion of a decidual membrane are the only diagnostic sign and symptoms. An inexperienced examiner is very apt to mistake a large serous or bloody cyst for uterine fibroma, and it is, indeed, often very difficult at first to distinguish the one from the other. But the uterine sound, carefully used, will show a great increase in the uterine cavity, when there is a fibroid, and a normal cavity in the case of the tubal affection.

The differential diagnosis between a large cyst of the tube and a fibrocystic tumor of the uterus is in some cases almost impossible; yet the increased size of the uterine cavity, as demonstrated by the use of the sound, will determine the question.

Special attention is directed to some points that are not dwelt on by authorities. Case of woman who suffered with pain in her pelvis. On examination uterus found retroverted and a mass on each side. It seemed as if the tubes were distended, so an operation was advised for the removal of the tubes. When the abdomen was opened instead of the tubes being distended they were normal in size. Both ovaries, however, gave indication of chronic ovaritis, and the supposed tubal affection proved to be a portion of the intestine occupying the sac of Douglas and containing two fæcal masses, occupying the position where distended tubes are usually found, overlapped by the uterus. Both ovaries were

removed, the adhesions separated, and the uterus returned to its normal position; but during her convalescence the patient was treated for ptosis, or prolapsus, of the bowel. William H. Skene (*Brooklyn Med. Jour.*, Aug., '98).

Prolapsed intestine and varicose veins have led the writer astray on several occasions. They no longer do so, for such patients are examined in the exaggerated Trendelenburg or exaggerated knee-chest position, when the tumor disappears. Placing the patient in either of these positions, a little pressure will raise the intestine sufficiently far up to know that it is not a tube, or it will sometimes slip out of the pelvis of its own accord. If there are many adhesions this will not take place, but the contents of the intestine can be pressed up out of the pelvis, so that the distended canal collapses and shows that it is not a tube. Placing the patient in this knee-chest position or the exaggerated Trendelenburg position, any present disappear, and the differential diagnosis is made in that way. A. J. C. Skene (*Brooklyn Med. Jour.*, Aug., '98).

GASTRO-ENTERITIS.

Treatment.—For infants suffering from gastro-enteritis due to milk infection, Dr. Wells advises absolute starvation for twelve hours,—no food of any kind being allowed during this period; cold boiled water, however, should be freely given, to which, if the child is very weak, from 10 to 15 drops of good whisky or brandy may be added. Copious irrigations of the lower bowel with boiled water containing a drachm of salt to the pint, are employed, one quart being used for each irrigation; these are best given through a catheter or small rectal tube. If there is much vomiting the stomach should be washed out with

sterilized boric-acid solution. It is well to give a small quantity of strychnine ($\frac{1}{120}$ grain) or other cardiac stimulant, hypodermically, just before lavage is performed. At the expiration of twelve hours, if the vomiting has ceased, and the stools look more natural, from $\frac{1}{2}$ drachm to a drachm of expressed beef-juice or liquid predigested food may be given every two hours. After that, if the infant has been bottle-fed, it is placed upon a modified milk diet. Breast-fed babies should be taken from the breast, and treated in all respects like those upon an artificial diet. The mother must be cautioned to bathe freely and before and after each nursing to wash the breasts and nipples with a strong solution of boric acid and water. Too much stress cannot be placed upon this injunction, since the mother's milk cannot fail to become infected the moment it reaches an unclean nipple. Charlotte C. West (*Phila. Polyclinic*, Aug. 6, '98).

GENU VALGUM.

The variety observed in the explanations given by different surgeons as to the origin of "knock-knee" is very puzzling to students, and moreover less misleading to practitioners. Experience will lead to recognition of various factors, which may be classified as follows: Changes in the soft parts,—for example, ligaments,—and softened and thickened epiphyseal cartilages; deformity in the lower end of the femur; deformity in the upper end of the tibia. These three factors are usually all present, but in some cases the deformity is the expression of only one of them; for instance, in a certain number of fairly-pronounced cases it was found that by supporting the internal condyle of the femur with one hand while the other pressed the

foot inwards toward the middle line of the body, the deformity could be made to disappear completely. In such cases the change of form is due to stretching of ligaments on the inner aspect of the joints, and to rachitic changes in the softened and thickened epiphyseal cartilages. Such cases never demand operative measures. The constant mention of operation in connection with genu valgum has a bad educational effect. Students come to regard operative treatment as a routine measure instead of one that is only exceptionally called for as a result of the neglect of sufficiently-early instrumental treatment. Clarke (*Brit. Med. Jour.*, Apr. 30, '98).

GERMAN MEASLES.

Diagnosis.—The characteristic enanthem observed in German measles is a macular, distinctly rose-red eruption upon the velum of the palate, the uvula, extending to, but not on to, the hard palate. The spots are arranged irregularly, not crescentically; are the size of large pin-heads; and are very little elevated above the level of the mucous membrane. The enanthem is very short lived, fading away within the first twenty-four hours. The claim that this enanthem is distinctive can be defended by comparison with the enanthem of those two diseases with which rubella is confounded. In scarlatina the enanthem appears from twelve to twenty-four hours before the eruption, on the pillars of the fauces in the form of the characteristic puncta, then rapidly spreads over the mouth in the form of a scarlet-red coalescing eruption, finally ending in desquamation, producing the strawberry-tongue, and lasting well into the second week of the disease. In measles the enanthem begins upon the soft palate from thirty-six to forty-eight

hours before the exanthem in the form of purplish or bluish papules, arranged crescentically, and extends over the cheeks, accompanied by the blue tongue. It is at its maximum with the beginning of the eruption, and may take as long as three or four days to disappear. Forchheimer (*Pediatrics*, July 1, '98).

GONORRHŒA.

Etiology.—Gonorrhœa may be innocently acquired. Experiments with pure culture of the gonococcus obtained from a gleet discharge of two years' standing gave the following results:—

1. Attempted reinfection of the original urethra with this culture was always a failure.

2. The culture, when transplanted to a coccus-free urethra, produced typical acute gonorrhœa.

3. Infection from this back again to the original urethra gave a fresh gonorrhœa, which, after a typical acute course of five or six weeks, again subsided to a chronic gleet. This most interesting experiment demonstrates that by passing an attenuated gonococcus through another subject—that is, through a fresh culture-ground—it becomes again virulent to a urethra which was immune to it before. This explains how an apparently-healthy man, if he have the gonococcus lurking anywhere in his urethral tract, may infect his hitherto-uninfected wife, and how he may be again infected from her. H. Brooks Wells (*New York Polyclinic*, May 15, '98).

Treatment.—In performing irrigation, it is well to get the patient used to having his anterior urethra thoroughly irrigated. When he finds he is not hurt, he will instinctively relax, and often without his knowledge some of the fluid finds its way into the bladder; then he

himself soon learns to relax the compressor muscle by trying to perform the act of urination. In this way the fluid flows into the bladder without any violence. The patient should sit on the edge of a chair, obtaining a comfortable position, sometimes by leaning back and relaxing himself completely. Then he holds a pus-basin in position with the left hand, while the physician sits on his right. In this way the bladder can often be filled rapidly, even when the height of the reservoir is but $2\frac{1}{2}$ feet. And it is important to avoid using any force whatever.

In some very few cases, after thoroughly cleansing the anterior urethra, the patient is placed on his back, and with a large hand-syringe, holding about 3 ounces, the bladder is filled from the meatus without the use of a catheter, after the method suggested by Guiard, of Paris. The strength of the solution used seldom exceeds 1 to 2000 for the anterior, and 1 to 4000 for the posterior, urethra; this is the best method to pursue in the majority of cases.

In addition to this treatment, argonin in 10-per-cent. solution may be used in the anterior urethra alone, injecting it by an ordinary urethral syringe and holding it in the urethra for a space of five to ten minutes.

In private work this should be done twice a day for three or four days according to the case, after that once a day; and in numerous cases the case comes to a complete close within a week, and incapable of returning except by a reinfection. The most brilliant cases are those seen early; but it may be applied at any stage. If it fails, it is because of infection of some para-urethral follicle which has been overlooked or some diverticulum within the urethra. In these early cases the disease seldom

reaches the posterior urethra, and often it is never necessary to irrigate the posterior urethra in a given case at all.

The course of the disease should be followed by microscopical examinations, the treatment carried on several days after the gonococci have disappeared, and the patient kept under observation for awhile. The only drugs given by the mouth are for the general condition or to keep the bowels open. Exercise, so far as possible, is interdicted. Alcoholics are absolutely forbidden during treatment, but attention is seldom paid to diet. The success of the treatment depends upon careful attention to details; so that it is never advocated giving the solutions to the patient to use himself. G. K. Swinburne (*Jour. Cutaneous and Genito-Urin. Dis.*, July, '98).

HYDROCEPHALUS, ACQUIRED.

Etiology.—It is probable that the most frequent causes of obstruction in cases of chronic hydrocephalus are simple fibrous closure of the foramen of Magendie, adhesion of the surfaces of the tonsils of the cerebellum to each other and to the margin of the fourth ventricle, and the presence of cysts between the arachnoid and pia at the postero-inferior aspect of the cerebellum.

Treatment.—A case of acquired hydrocephalus was operated on and drainage established through the fourth ventricle. The trephine was applied to the occipital bone in the mesial line a little above the foramen magnum; although the skull is particularly thick at this point, and the sinus in the falx cerebelli requires to be ligated, this is the easiest and most satisfactory approach to the fourth ventricle. The accumulation of cerebrospinal fluid in this case was due to adhesions between the two tonsils of

the cerebellum and the sides of the medulla, the separation of which was followed by the escape of the imprisoned fluid. In the subsequent course of the case a large quantity of cerebrospinal fluid escaped daily from the wound. The operation is one that should be given trial in cases of chronic basilar meningitis of both the tuberculous and non-tuberculous varieties. Bruce and Stiles (*Scottish Med. and Surg. Jour.*, March, '98).

HYSTERIA AND PELVIC DISEASE.

The psychical symptoms of hysteria are important to the gynæcologist. The patient is, as a rule, exceedingly impressionable. She is open to suggestion, especially as regards her pelvic condition. Hysteria is a psychoneurosis because of the prominence of these psychical manifestations. Neurasthenia may exist without pelvic disease. If both co-exist they have no relation with each other. If pelvic disease exists with neurasthenia the pelvic symptoms become more marked because of increased irritability. Hysteria may exist without pelvic disease. The possibility is denied of nervous and mental disease arising from pelvic operations. The pelvic condition should be operated on for the local condition only, and not to relieve the nervous condition. In cases of profound hysteria an operation should never be undertaken unless the surgical condition is very urgent. The hysteria should first be cured in order to prevent the disastrous effects of the operation upon the nervous condition. The insanities are not due to local organic disease, but to disease of the neurons as a result of various derangements of tissue-metabolism. Pelvic operations will not cure insanity. F. X. Dercum (*Med. News*, June 25, '98).

INTESTINAL PERFORATION.

Diagnosis.—Ether as a means of diagnosis of intestinal perforations is more practical and better than hydrogen-gas, since it is always at hand, and necessitates no special apparatus, aside from what is already found in the surgeon's possession, and further, no objections can be given to its use on account of noxious properties, unfounded as these objections may be as to the hydrogen-gas. The quantity of ether being infinitesimal (1 drachm or less being sufficient to give up the necessary vapor, which is diluted many, many times by the air that carries it) absolute freedom from anaesthesia is assured. The method is simple: A small quantity of ether is poured into the bottle accompanying the aspirator of so-called French pattern. A soft-rubber catheter or rectal tube (preferably a long, glass douche-tube connected by rubber tubing) is jointed to one of the pipes of the aspirator-bottle. The air-pump is joined to the other pipe, and air forced into the bottle becomes mixed with ether-vapor and passes on through the alimentary canal, the rectal tube being inserted as far as necessary. This mixture, which may be called ether-air, rapidly finds its way through the coils of intestines, giving forth strong rumblings as it progresses onward, causing distension on entering the stomach, from whence it is belched, provided no perforation exists along its pathway. Upon reaching a perforation, however, the ether-air escapes into the peritoneal cavity, tympanites more rapidly develops, and upon dilating the wound-entrance down to the peritoneum, quickly comes out into the world, being at once recognized by its odor and the hissing sound of escaping gas. Upon opening the abdomen the distension of the intestine from the rectum up can

be followed to the point of injury where the same odor and hissing are noted and the wound repaired. Continuing the search till no more ether-air is found escaping from the intestines one may rely upon its efficacy. E. M. Sutton (*Jour. Amer. Med. Assoc.*, July 23, '98).

KRYOPHINE.

Kryophine is produced by heating p-phenetidin with methoxyacetic acid to 248°-266° F.; it is tasteless and odorless; soluble in 600 parts of cold, 52 parts of boiling water; also in alcohol, ether, chloroform, glycerin, and fixed oils. Under its influence the pulse becomes fuller and stronger, with a disappearance of diastole. Abnormal temperature is reduced surely and promptly, in a marked degree, and extending over several hours of time, without rigor or depression, and rarely attended by diaphoresis. Blood-pressure is increased. Respiration is not affected. It is eliminated in about six hours, by the kidneys principally, although it does not affect diuresis, and may be found in the urine fifteen minutes after its ingestion. The rapid and more or less complete saponification by the stomach and intestinal juices is probably the secret of its power and the reason for its prompt action. It controls neuralgic pain in a marked and sometimes almost magical manner, and in some persons produces a tendency to sleep. For the average adult 8 grains seems to be the dose giving best results, and is best given in powdered form, dry, upon the tongue. The tablets are not advisable until first pulverized. John H. Curtis (*Therapeutic Gazette*, May, '98).

MEASLES.

Diagnosis.—The Koplik spots are round, slightly raised, bluish-white efflorescences, having minute red centres.

They measure from 0.01 to 0.03 inch in diameter and are situated upon the inner surfaces of the cheeks, less often upon the mucous membrane of the lips, and rarely (in one instance) upon the tongue. They may occur on one or both sides, most often in the vicinity of the back teeth, and the usual number is from six to twenty, though hundreds may be present. They may be seen in daylight, or by a strong incandescent light, but not by lamplight. These spots seldom run together and cannot be rubbed off. They usually make their appearance on the first or second day of the prodromal stage and increase in number until the skin eruption appears, when they remain without change for three or four days and then disappear. Slawyk (*Deut. med. Woch.*, Apr. 28, '98).

MILK AS A CULTURE-MEDIUM.

The composition of milk makes it a nutritive medium for every germ whose biology is at present understood. An investigation carried on in Boston showed that when healthy cows, in a clean place, were carefully milked into a sterilized flask, the milk contained only 530 bacteria per cubic centimetre, but when the ordinary milk-pail is used and the milk is conducted in the common way, there were, on an average, 30,500 bacteria per cubic centimetre immediately after milking. It should be remembered that the germs of diphtheria, cholera, typhoid fever, and scarlet fever develop at ordinary room-temperature, and that such growth does not affect the gross appearance of the milk. J. W. Strickler (*Phila. Med. Jour.*, July 9, '98).

MILK: ITS ABSORPTION vs. ITS DIGESTION.

Milk alone is sufficient to meet the needs of the body. Intravenous injection

tion of fresh milk has been safely practiced, showing that the economy can assimilate it without previous digestion and absorption. Working on this theory, the author has adopted the following plan for the rapid absorption of milk without previous curdling and digestion. A number of hours after a meal (usually three or four) the food disappears from the stomach, with all gastric juice, and the mucous surface becomes alkaline. This is the "alkaline tide" of the stomach. If at this time milk, free from fat, fresh and alkaline, and at the temperature of the body, be taken, it will excite no secretion of gastric or pancreatic juice on account of its freedom from all irritant qualities, and it will, therefore, pass at once in an unaltered state into the absorbents and system. This saves nature much work and avoids the disturbances of coagulation. It enters the blood-current more quickly, and in no way disturbs the appetite for regular meals, even increasing the latter. Milk can thus be taken by patients who cannot take it with their meals. Bulkley (*Phila. Med. Jour.*, July 2, '98).

MYXŒDEMA IN THE NEGRO.

Symptoms.—The existence of myxœdema among the black races has been universally denied, but there are myxœdemic conditions sometimes seen that exhibit a peculiar thickening of the skin, local in distribution, identical in all respects with that present in cases of sporadic cretinism in the Caucasian race, but less diffuse in character. In 7 cases the thyroid gland was either not palpable, or below normal in size; in one case it was enlarged. The hair showed alteration in two cases, being coarse, thin and rough. The bones were abnormally broad in one case; the secretions of the skin were normal; even over the myxœ-

dematous areas there was little of the dryness and roughness usually characteristic of the disease. The features were broadened and rendered coarse in two cases, the abdomen pendulous in one. The blood was carefully examined, but no abnormalities were found in the shape or diameter of the red corpuscles in any case.

Treatment.—The excretion of urea rose to about the normal in nearly all under thyroid administration, while previously it had been below the normal. In four cases the administration of dry thyroid gland removed the jelly-like thickenings of the skin, proving fairly conclusive that the swellings were of the same general nature as myxœdema; and all the cases improved mentally during the period of administration. Berkley (*Amer. Jour. Insan.*, April, '98).

NEURASTHENIA.

Etiology.—The disorders of woman's pelvic organs have no more to do with her nervous and mental diseases than lesions elsewhere in her body; indeed, they have less to do with her psychoses and neuroses than most of her other organs, for, as in the male sex, the chief causes of their neuropsychoses are to be sought in intrinsic disorders of the nervous system itself, or in perverted nutrition of the nervous system dependent upon affections of the gastro-intestinal tract, kidneys, liver, lungs, heart, etc., and upon pathological blood states. It is true that puberty, adolescence, the puerperium, menstruation, and the menopause are often closely related to the outbreak or to the exacerbation of many nervous and mental disorders, but the pelvic organs themselves play but a small rôle in these physiological commotions. They have to do with the whole organism of woman. It is not denied

that pelvic diseases in women attended by exhausting pain may give rise to neurasthenic and hysterical states, but the influence of exhausting pain in these organs is no greater than similar exhausting pain elsewhere in the body. Nor is it denied that disorders of the female organs which affect the nutrition of the nervous system, such as excessive hæmorrhage, or suppurative processes, may be important factors in inducing functional neurosis and even insanity, though disordered blood states brought about by pelvic disease are very infrequent as compared with disordered blood states dependent upon disease elsewhere. There is no evidence whatever to support the opinion that insanity was ever due to a mere reflex influence from pelvic disease. In insanity the two great etiological factors are hereditary instability and some physical or moral stress directly affecting the nervous system. There has never been brought forward any evidence whatever to show that either epilepsy or chorea can be induced by disease of the female organs. Frederick Peterson (*Annals of Gynecology and Pediatrics*, Aug., '98).

OPHTHALMIA NEONATORUM.

In the twenty-five largest cities of the United States the proportion of the blind is, with two exceptions, smaller than in the States in which these cities are situated, or, taking all the cities of more than 50,000 inhabitants together, about 33 per cent. less blind are found in them than the average for the entire country. When the different factors in the production of blindness are examined, whether congenital or acquired, or, under the latter class, whether due to traumatism, general disease, or to local disease, these factors are all practically the same, or are made equal in

city and country, with one exception, namely, ophthalmia of infancy. A rather extended inquiry concerning the habitual practice of physicians in country almshouses, in hospitals and elsewhere in the State of New York indicates that more attention is given to guarding against ophthalmia of infancy in the cities than in the country. The tendency to the habitual neglect of prophylaxis tends to make a rapid difference in the distribution of the blind, estimated at possibly 14 to 1. It is at least the most apparent cause of this difference and probably accounts for the greater part of it. It follows from this apparently warrantable conclusion that if as great care were taken in general throughout the country as is given on the average in the cities to prophylaxis the number of blind in the United States would be decreased in a single generation by several thousands. Lucien Howe (*Med. News*, Aug. 6, '98).

PEMPHIGUS.

Treatment.—In the treatment of pemphigus vulgaris no reliance can be placed on the specifics; arsenic and quinine with tonics can be used with some hope of doing good. The arsenic should be used in the form of Fowler's solution, commencing with small doses and gradually increasing to tolerance. Quinine should be given in large doses, 20 to 30 grains every second or third day. Tonics in general should be given, such as cod-liver-oil, malt, iron, strychnine, etc., to improve the lowered vitality of the patient. Local treatment is of great importance; constant baths with the use of mild ointment, with vaselin as a base, and bismuth subnitrate, aristol, and oxide of zinc as constituents. In a disease like pemphigus, where the patient is more or less flayed, the hot bath

offers a medium in which the patient can live without such anguish as constantly tortures him, and with a greater chance of recovery. Very little true absorption of the water or the substance contained in it takes place in the bath, but a certain amount of the imbibition of the water and substances contained therein may occur. The baths should be given every day, or two or three times daily, if necessary to give the patient comfort; after the bath he should be anointed with a mild paste, of the formula before mentioned, enveloped in absorbent cotton, same being held in place by roller bandages. The temperature of the bath should be about 90° F., the patient should be immersed from one-half to two hours, and plenty of Castile soap used, so that the superficial layers of the epidermis should be cleansed, thereby favoring imbibition. The temperature of the bath should be kept up by the addition of hot water, and the room in which the patient takes his bath kept at an even warmth. A patient should not be put into a bath

after a hearty meal; the best time to give the bath is in the afternoon between 3 and 6 o'clock. Secretion from the bullæ should be tested, and if it is found to be neutral, simple hot bath to be given; if acid, an alkaline bath, made so by the addition of 4 ounces of bicarbonate of soda to 30 gallons of water. If alkaline, an acid bath is given, made so by the addition of 1 1/2 ounces of nitric acid and 1 ounce of hydrochloric acid to 30 gallons of water. The daily diet should be highly nourishing; general hygienic measures and good ventilation should be observed. Formaldehyde fumigation is followed by success; two burners generally used twice a day; this is as long as can be comfortably borne by the patient. They soon become accustomed to its use; the first day they will allow the lamp to burn about ten minutes; by the end of the first week one to two hours. The large-size bullæ had best be opened at the most dependent portion and the fluid allowed to escape. R. P. Izlar (Georgia Jour. of Med. and Surg., Aug., '98).

New Books and Monographs Received.

The editor begs to acknowledge with thanks the receipt of the following books and monographs :—

Does the Theory that Typhoid Fever can be Aborted Conflict with Any Established Law of Pathology or with Any Known Scientific Fact? By John Eliot Woodbridge, M.D., Cleveland, Ohio, 1898.—Renal Suppuration, Catarrhal, Specific, and Traumatic, and the value of Micro-urinalysis of the Urinary Sediment as an Aid to Definite Diagnosis of It. By Thomas H. Manley, M.D., New York City, 1897.—A Clinical Study of Kryofine. By Sidney V. Haas, M.D., and J. Bennett Morrison, M.D., New York City, 1898. The Production and Sale of Antitoxin by the New York Board of Health. By A. M. Phelps, M.D., New York, 1898.—An Exhibition of Radiographs, with Remarks. By A. V. L. Brokaw, M.D., St. Louis, Mo., 1897.—Some Conclusions Drawn from Experiences in Pelvic Surgery. By A. V. L. Brokaw, M.D., St. Louis, Mo.—Transactions of the Section on Ophthalmology: College of Physicians of Philadelphia. March 15 and April 19, 1898.—Forest-growth and Sheep-grazing in the Cascade Mountains of Oregon. By Frederick V. Colville. U. S. Department of Agriculture, 1898.—

Our Trade with Spain, 1888-1897. Prepared by Frank H. Hitchcock. U. S. Department of Agriculture, 1898.—The San Jose Scale in 1896-1897. By L. O. Howard. U. S. Department of Agriculture, 1898.—Spain's Foreign Trade. Prepared by Frank H. Hitchcock. U. S. Department of Agriculture, 1898.—The Inspection of Meats for Animal Parasites. Prepared under the direction of Dr. D. E. Salmon. U. S. Department of Agriculture, 1898.—Final Report of the Crops of 1897. U. S. Department of Agriculture, 1898.—Report of the Public Education Association of Philadelphia Made at the Seventeenth Annual Meeting, 1898.—The Grain-smuts: How They are Caused and How to Prevent Them. By Walter T. Swingle. U. S. Department of Agriculture, 1898.—Cattle-ranges of the Southwest. By H. L. Bentley. U. S. Department of Agriculture, 1898.—Epidemic Cerebro-spinal Meningitis and its Relation to the Other Forms of Meningitis. A Report of the State Board of Health of Massachusetts, 1898.—Report of the Health Department of the City and County of San Francisco for the Fiscal Year Ending June 30, 1897.—Farmakodynamiska Studier A Det Isolerade Och Offerlevande Däggdjurshjartat af Karl Hedbom. Upsala, 1896.—Ueber Giftige Eiweisse Welche Blutkörperchen Verkleben. Von M. Elfstrand, Upsala, 1897.—Arsberättelse Fran Sabbatsbergs Sjukhus. 1 Stockholm för 1895-96. By Dr. F. W. Warfvinge, Stockholm, 1898.—A proposito della cura dell'ozena col siero antidifterico, 1898.—Discorso del Prof. Vittorio Grazzi all a inaugurazione del Terzo Congresso della Societa Italiana do Laringologia, Otologia, e Rinologia, avvenuta nell a R. Università di Roma la mattina del 28 Ottobre, 1897.—Di Una Grave Complicazione Avvenuta Dopo l'Asportazione di un Papilloma della Laringe per il Prof. Vittorio Grazzi, Florence, 1897.—Note Oto-rino-laringologiche. By Prof. Vittorio Grazzi, Florence, 1898.—Sur le Traitement des Sinusites. By E. J. Moure, Bordeaux, 1897.—Sur un Cas de Surdite Complete a la Suite d'une Meningite Aigue par Diplococcus de Fraenkel et presentation de la Malade. By Prof. Vittorio Grazzi, Florence, 1897.—Post-febrile Insanity. By Alexander L. Hodgdon, M.D., Baltimore, 1897.—Notes on the Pathology and Bacteriology of Appendicitis. By Chas. F. Craig, M.D., Danbury, Conn., 1897.—The Bacteriology of Epidemic Carotitis, with Description of a Diplo-Bacillus Found in the Blood and Urine. By Chas. F. Craig, M.D., Danbury, Conn., 1898.—The Branched Form of the Bacillus Tuberculosis in Sputum. By Chas. F. Craig, M.D., Danbury, Conn., 1898.—The Centrifuge as an Aid in Diagnosis. By Chas. F. Craig, M.D., Danbury, Conn., 1898.—Sanitarium Treatment of Pulmonary Tuberculosis. By J. Edward Stubbart, M.D., Liberty, N. Y., 1898.—The First Recognized Case of Yellow Fever in Mobile in 1897, with Comments and Deductions. By Edwin L. Maréchal, M.D., Mobile, Ala., 1898.—The Treatment of Diphtheria with Diphtheria Antitoxin. By Edwin Rosenthal, M.D., Philadelphia, Pa., 1895.—The Influence of Antitoxine in the Treatment of Laryngeal Diphtheria With and Without Intubation. By Edwin Rosenthal, Philadelphia, Pa., 1898.—Injuries to the New Born in Cross, Complex and Breech Presentation. By Edwin Rosenthal, M.D., Philadelphia, Pa., 1898.—Uterine Moles. By Edwin Rosenthal, M.D., Philadelphia, Pa., 1898.—Serum Therapy in Diphtheria. By Edwin Rosenthal, M.D., Philadelphia, Pa., 1896.—Etiology, Prevalence and Treatment of Hysteria. By U. O. B. Wingate, M.D., M.M.S.S., Milwaukee, 1898.—Physical Characteristics of Ten Thousand Men. By Lt.-Col. Charles Adams, Chicago, Ill., 1898.—The Essential of the Art of Medicine. By J. H. Musser, M.D., Philadelphia, Pa., 1898.—The Diagnostic Importance of Fever in Late Syphilis. By J. H. Musser, M.D., Philadelphia, Pa., 1892.—Renal Calculus. By J. H. Musser, M.D., Philadelphia, Pa., 1898.—Symposium on the Pathology of the Diseases of the Cardio-Vascular System. By J. H. Musser, M.D., and J. D. Steele, M.D., Philadelphia, Pa., 1898.—The Prevention of Diseases now Preying upon the Medical Profession. By Leartus Connor, A.M., M.D., Detroit, Mich., 1898.—Address of the President. By W. Murray Weidman, M.D., Reading, Pa., 1898.—A


Report upon the Grasses and Forage Plants and Forage Conditions of the Eastern Rocky Mountain Region. By Thomas A. Williams, "U. S. Department of Agriculture," 1898.—Yellow Fever: Its Nature, Diagnosis, Treatment, and Prophylaxis, and Quarantine Regulations Relating Thereto. By Officers of the U. S. Marine Hospital Service. Prepared under Direction of the Supervising Surgeon-General, 1898.

EDITORIAL STAFF.

ASSOCIATE EDITORS.

(List Revised August 1, 1898.)

J. GEORGE ADAMI, M.D., MONTREAL, P. Q.	S. G. GANT, M.D., KANSAS CITY, MO.	WILLIAM OSLER, M.D., BALTIMORE, MD.
LEWIS H. ADLER, M.D., PHILADELPHIA.	J. McFADDEN GASTON, M.D., ATLANTA, GA.	F. A. PACKARD, M.D., PHILADELPHIA.
JAMES M. ANDERS, M.D., LL.D., PHILADELPHIA.	J. E. GRAHAM, M.D., TORONTO, ONT.	LEWIS S. PILCHER, M.D., BROOKLYN, N. Y.
G. APOSTOLI, M.D., PARIS, FRANCE.	JULES GRAND, M.D., PARIS, FRANCE.	WILLIAM CAMPBELL POSEY, M.D., PHILADELPHIA.
A. D. BLACKADER, M.D., MONTREAL, P. Q.	EGBERT H. GRANDIN, M.D., NEW YORK CITY.	W. B. PRITCHARD, M.D., NEW YORK CITY.
E. D. BONDURANT, M.D., MOBILE, ALA.	LONDON CARTER GRAY, M.D., NEW YORK CITY.	JAMES J. PUTNAM, M.D., BOSTON.
DAVID BOVAIRD, M.D., NEW YORK CITY.	J. P. CROZER GRIFFITH, M.D., PHILADELPHIA.	GEORGE H. ROHÉ, M.D., SYKESVILLE, MD.
L. BROCC, M.D., PARIS, FRANCE.	A. GOUGUENHEIM, M.D., PARIS, FRANCE.	ALFRED RUBINO, M.D., NAPLES, ITALY.
WILLIAM BROWNING, M.D., BROOKLYN, N. Y.	C. M. HAY, M.D., PHILADELPHIA.	LEWIS A. SAYRE, M.D., NEW YORK CITY.
WILLIAM T. BULL, M.D., NEW YORK CITY.	FREDERICK P. HENRY, M.D., PHILADELPHIA.	REGINALD H. SAYRE, M.D., NEW YORK CITY.
CHARLES W. BURR, M.D., PHILADELPHIA.	EDWARD JACKSON, M.D., DENVER, COL.	SOLOMON SOLIS-COHEN, M.D., PHILADELPHIA.
DUDLEY W. BUXTON, M.D., M.R.C.P. LONDON, ENGLAND.	NORMAN KERR, M.D., F.L.S., LONDON, ENGLAND.	H. W. STELWAGON, M.D., PHILADELPHIA.
HENRY T. BYFORD, M.D., CHICAGO, ILL.	EDWARD L. KEYES, JR., M.D., NEW YORK CITY.	D. D. STEWART, M.D., PHILADELPHIA.
J. ABBOTT CANTRELL, M.D., PHILADELPHIA.	H. KRAUSE, M.D., BERLIN, GERMANY.	LEWIS A. STIMSON, M.D., NEW YORK CITY.
WILLIAM B. COLEY, M.D., NEW YORK CITY.	E. LANDOLT, M.D., PARIS, FRANCE.	G. ARCHIE STOCKWELL, M.D., NEW YORK CITY.
P. S. CONNER, M.D., LL.D., CINCINNATI, OHIO.	ERNEST LAPLACE, M.D., LL.D., PHILADELPHIA.	B. J. STOKVIS, M.D., AMSTERDAM, HOLLAND.
FLOYD M. CRANDALL, M.D., NEW YORK CITY.	R. LÉPINE, M.D., LYONS, FRANCE.	LOUIS McLANE TIFFANY, M.D., BALTIMORE, MD.
ANDREW F. CURRIER, M.D., NEW YORK CITY.	F. LEVISON, M.D., COPENHAGEN, DENMARK.	CHARLES S. TURNBULL, M.D., PHILADELPHIA.
JUDSON DALAND, M.D., PHILADELPHIA.	A. LUTAUD, M.D., PARIS, FRANCE.	HERMAN F. VICKERY, M.D., BOSTON, MASS.
N. S. DAVIS, M.D., CHICAGO, ILL.	F. MASSEI, M.D., NAPLES, ITALY.	RIDGELY B. WARFIELD, M.D., BALTIMORE, MD.
F. EKLUND, M.D., STOCKHOLM, SWEDEN.	E. E. MONTGOMERY, M.D., PHILADELPHIA.	F. E. WAXHAM, M.D., DENVER, COL.
AUGUSTUS A. ESHNER, M.D., PHILADELPHIA.	JULES MOREL, M.D., GHENT, BELGIUM.	J. WILLIAM WHITE, M.D., PHILADELPHIA.
J. T. ESKRIDGE, M.D., DENVER, COL.	HOLGER MYGIND, M.D., COPENHAGEN, DENMARK.	W. NORTON WHITNEY, M.D., TOKIO, JAPAN.
CHRISTIAN FENGER, M.D., CHICAGO, ILL.	W. P. NORTHRUP, M.D., NEW YORK CITY.	JAMES C. WILSON, M.D., PHILADELPHIA.
SIMON FLEXNER, M.D., BALTIMORE, MD.	H. OBERSTEINER, M.D., VIENNA, AUSTRIA.	C. SUMNER WITHERSTONE, M.D., PHILADELPHIA.
LEONARD FREEMAN, M.D., DENVER, COL.	CHARLES A. OLIVER, M.D., PHILADELPHIA.	WALTER WYMAN, M.D., WASHINGTON, D. C.



FORMULA:
 B Ferri Sulph. 1½ grs.
 Potass. Carb. 1½ grs.
 Dose—1-2.
 Price, 40c. per 100.

The best method for prescribing Iron

As indicated
in

ANEMIA
CHLOROSIS
PHTHISIS
etc., etc.

Pil. Chalybeate

Offers ferrum in the
EASILY ASSIMILABLE FORM OF
FERROUS CARBONATE.

PROF. NIEMEYER writes: "For more than twenty years I have used these pills almost exclusively in Chlorosis, Phthisis, etc., and have witnessed such brilliant results from them in a large number of cases that I never experimented with other preparations of iron."

PIL. CHALYBEATE COMP.


Is same formula as Pil. Chalybeate with ¼ gr. Nux Vomica added to increase the tonic effect.

Please specify

WM. R. WARNER & CO.

PHILADELPHIA. NEW YORK.

CHICAGO.



SCHERING'S PIPERAZINE WATER

is the most approved form for administration of Piperazine. Its efficacy in the treatment of RHEUMATISM, GOUT, and its solvent power upon Uric Acid Concretions in the Joints and Stone in the Bladder, have been attested to by Biesenthal, Ebstein, Heubach, Meisels, David D. Stewart, Hildebrandt, Schweninger, Wilcox, and many others.

In a recent contribution to the *Medical News*, November 27, 1897, by Prof. R. W. WILCOX: "A Phase of the Treatment of Goutiness," the author arrives at the following conclusions:

"That Piperazine, administered in the form described (Piperazine Water) in this paper, is the remedy of choice for the elimination of uric acid, not only in this (neurotic lithemia) but in other pathologic conditions dependent upon the same cause."

PIPERAZINE WATER IS SUPPLIED
IN CHAMPAGNE QUARTS.
DAILY DOSE ONE BOTTLE.

Excerpts from Literature Sent on Demand by

Lehn & Fink, Manufacturing Chemists, New York.

In writing to advertisers mention Monthly Cyclopædia

NOTES AND COMMENTS.

THE F. A. DAVIS COMPANY, Publishers.

GEO. B. CREVELING, Business Manager.

General Greely's 278 Days of Death.—

The true story of those 278 days of suffering by Greely's heroic little band of explorers in the Arctic region has been told by General Greely himself, for the first time, for the October Ladies' Home Journal. For years General Greely has kept an unbroken silence about his fearful experience and that of his companions, as they dropped dead one by one at his side, and it was only after the greatest persuasion that the famous explorer was induced to write the story.

The Battle-Creek Sanitarium.—

This institution, under the able superintendence of Dr. J. H. Kellogg, is known throughout the country for the completeness of its equipment as well as the numerous scientific and clinical investigations which are carried out within its walls. It has the advantage over many hospitals in being amply provided with conveniences for the utilization of physiological in addition to the ordinary medicinal remedies. The location and surroundings of the sanitarium are extremely pleasant.

Wet Dressings in Surgery.—

From an interesting paper of the above title presented to the section on Surgery and Anatomy at the Denver meeting of the American Medical Association we reproduce the following passages:—

"It is clear that to carry out the idea of a practical dressing it must be:—

"1. Antiseptic. This applies not only to the effect of the application to the part, but to the corporate substance itself.

"2. Permanent. This is necessary in order to avoid too frequent removals as well as to preserve itself from deterioration.

"3. Non-irritating. There is nothing more delicate, more easily disturbed than the formative principles of tissue; so that care must be taken in the dressing of all lesions of surface lest the agents used should arrest the tender process of repair, as well as protect it from the invasion of destructive germs.

"4. Constructive. While the majority of lesions, especially those of traumatic origin, if not interfered with by destructive germs will heal rapidly of themselves, there are very many which require not only this negative condition, but also a positive stimulation of function in the constructive elements of the part."

The fulfillment of these conditions has been the aim of the surgical pharmacist, but it was to Sir Astley Cooper that the credit is due for his foreseeing therapy leaping over, as it were, the dark chasm which separated the triumphs of his surgical genius from the science-illuminated land of modern surgical pathology. It was he who, without the knowledge of the bacteriological factor in the great problem of surgical treatment, by the intuition of genius gave to us the essential principles of external dressing for surface-lesions. His formula, however, was open to the objection of violating one of the conditions herein laid down—that of permanence—in that lard was used instead of petrolatum, which has been since discovered and is now substituted in the

In writing to advertisers mention *Monthly Cyclopædia*

Rx: Ung't. Resinol $\mathfrak{z}\text{i}$ — iii .

THOROUGHLY EFFICIENT AS AN
ANTIPRURITIC, ANTISEPTIC,
LOCAL ANTIPHLOGISTIC & SKIN NUTRIENT

And is now being successfully used and prescribed by over 40,000 PHYSICIANS in the United States and Canada

For Eczema, Erysipelas, Pruritus and all Itching, and
Inflammatory Skin Diseases.

Rx: Elix. Cascanata $\mathfrak{z}\text{iv}$ — x .

LAXATIVE, ALTERATIVE, ANTACID AND TONIC.

Stimulates the elimination of effete matter and restores the
normal alkalinity of the blood.

A specific for habitual constipation and the digestive derangements resulting therefrom.

RESINOL CHEMICAL CO., = Baltimore, Md.

Lactopeptine
contains
all known
substances employed by
nature in the diges-
tion of all kinds
of food.
The N.Y. Pharmaceutical
Association
Yonkers, N.Y.

Notes and Comments (continued).

preparation known as unguentine, which is an ideal formula constructed along the lines of that suggested by Sir Astley Cooper, but altered to the conditions of modern aseptic surgery. The irritating effects of the ordinary alum has also in some way been obviated, furnishing thus a typical dressing for surface-lesions. We are free to admit that for all external dressings we have found the highest fulfillment of modern aseptic or antiseptic surgery in the preparation just mentioned.

The Blood is the Life.—The purity of the circulating fluid is the prime necessity of normal and healthy nutrition. Sooner or later, more or less severely, the different organs and tissues of the body suffer when the blood is impoverished or contaminated. In anæmia, in many forms of wasting and inflammatory diseases and in those which diminish assimilation experience has taught the benefit of supplying blood itself directly to the system. This is accomplished by means of bovine, which is pure and unaltered blood from the arteries of the bullock. Bovine can be administered by the mouth, rectum, or subcutaneous injection. Sprayed upon the mucous membrane and followed by hydrogen dioxide it is serviceable in catarrhal diseases of the air-passages and diphtheria.

Ye Surgeons.—Judging from frequent reports in current medical literature Lysol is being extensively used by many of our well-known surgeons. The saponaceous character of its solution is very advantageous in many cases, rendering a special surgical soap unnecessary in preparation for operations. It is a non-toxic antiseptic, soluble in water, and is

especially used in gynæcology for intra-uterine irrigation after curettage (Prof. Augustin H. Goelet, School of Clinical Medicine, New York), and is not caustic like carbolic acid and preferred to it as well as to bichloride of mercury for vaginal douches in obstetrics (Dr. Mary H. McLean), and for cleaning lacerated wounds and contusions (Dr. E. O. Plumbé, "Railway-surgeon," April, 1898). It is used in $\frac{1}{2}$ per cent. to 1 per cent. solutions in water. ("Notes on New Remedies.")

Epilepsy.—The Tilden Co., St. Louis, Mo. Dear Sirs:—Inclosed find post-office money-order for \$1.00. Please send by return mail one ounce tablets of hydrocyanate of iron and valerian. My case of epilepsy is doing fine, has had no trouble for the past three months, and appears as well as he ever was. Works hard every day on hay-bales, tying and lifting heavy bales of hay. Will send full history of case when we get through if you desire it. Respectfully, F. E. Wilson, Westminster, Cal., August 11, 1898.

"A New Created World."—If you leave home for the summer, go where you will find almost literally a new world. Such a spot is Yellowstone Park. There is no place like it in the world. You will be astonished at what you see and learn there. A month's outing in those mountains 7000 feet above the sea-level, in an atmosphere pure and invigorating, would add years to your life. There are four modern hotels, good roads—maintained by the United States Government—for riding, driving or cycling; riding-horses and pack-trains. Geysers, canyons, mud-springs, lakes, cataracts, and cascades in large number are crowded into a space 54 by 62 miles

In writing to advertisers mention Monthly Cyclopædia

SPRING TRIPS! . . .

Clyde Line Excursions,
Including Meals and Accommodations,

New York to

Charleston, S. C.,

and return

Intermediate Round Trip, \$24.00;

First-Class Round Trip, \$32.00.

New York to

FLORIDA

and return

Intermediate Round Trip, \$35.30;

First-Class Round Trip, \$43.30.

\$3.75 additional includes round trip transportation on the beautiful

ST. JOHN'S RIVER.

For full particulars see your nearest ticket agent, or write to

PASSENGER DEPARTMENT:

201 Washington Street, Boston, Mass.,
12 S. Delaware Ave., Philadelphia, Pa.,
5 Bowling Green, N. Y.

WM. P. CLYDE & CO., General Agents.

T. G. EGER, Traffic Manager.

NORTH & SOUTH
TRAVEL BY THE
CLYDE LINE



NEW YORK
CHARLESTON, S.C.
AND
FLORIDA

UNSURPASSED
PASSENGER
ACCOMMODATIONS

YOU'VE SEEN
NEW YORK & JACKSONVILLE FLA.
WITHOUT LEAVING

WM. P. CLYDE & CO., GENERAL AGENTS.
5 BOWLING GREEN, NEW YORK. 12 SOUTH DELAWARE AVE. PHILA.

↓
ESPERANZA (DRY).



↑
HYGEIA (SWEET).

Unfermented Concord Grape Juice.

FROM CHOICE, SELECTED GRAPES. NO
ALCOHOL. NO ANTISEPTICS.

Nature's Best Drink and Tonic.

FOR THE SICK. FOR THE WELL.

Prepared by

SNOW GRAPE JUICE CO.,

PENN YAN, N. Y.

SEND 30 CENTS IN STAMPS FOR PINT BOTTLE.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

in size. If you can afford it, you wrong yourself by not going there, especially if you are a patriotic American. If you cannot afford it, then go to beautiful Leech Lake in Minnesota, a new and popular resort. New hotels, fine fishing and hunting, and good boating at reasonable rates. Send Charles S. Fee, of Northern Pacific Railway, St. Paul, Minn., six cents for Wonderland, '98, that describes by pen and picture both resorts. Mention where advertisement was seen.

Validol.—The above is the title of a new preparation introduced by Dr. George Schwersenski, of Berlin, who has described its properties and uses in an article published in the *Therapeutische Monatshefte*. Validol is a chemically-pure combination of menthol and valerianic acid. It contains a proportion of free menthol, which adds to the stimulant effect. Validol is a clear, colorless liquid with a consistence similar to that of glycerin. It has an agreeable odor and a cool, slightly-bitter taste. This fluid will hold any amount of menthol in solution, and therefore the latter can be administered in any required dose. Validol, containing 30 per cent. of free menthol, answers most requirements. As a rule, 10 to 15 drops of

validol given daily once or oftener according to the severity of the case will be sufficient. This preparation has proved an excellent remedy in hysteria and neurasthenia. As a stomachic validol removes nausea and malaise and promotes the appetite. It seems to be useful also as a carminative.

Prof. Gennaro Scognamiglio, of Naples, has given validol with much satisfaction in seasickness, in which he considers it to have both specific and prophylactic properties.

Validol is manufactured by the Vereinigte Chininfabriken Zimmer and Co., of Frankfort-on-the-Main, Germany (and sold by R. W. Phair & Co., New York City).

The Omaha Exposition.—When the

World's Fair at Chicago ceased to exist, it was supposed that we should ne'er look upon its like again. However, the Trans-Mississippi Exposition at Omaha has effectively reproduced in similarity all of the buildings which made the White City so attractive in 1893.

It does not now take weeks to wander through grounds and structures and then be compelled to go away with a jumble of ideas, for the Omaha Exposition people have profited by past experience, and have so improved the arrangement



The Trained Nurse Knows,
Platts' Chlorides.
The True Disinfectant.

Platts' Chlorides, **The True Disinfectant.**

An odorless, colorless liquid; powerful, safe, and cheap; indorsed by over 23,000 physicians; sold everywhere in quart bottles only. Prepared only by

HENRY B. PLATT,

Platt Street, New York.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

of exhibits that no more than two or three days of time need be consumed in admiration and inspection of the marvelous resources of the West, collected together in the chief city of Nebraska.

Even the new Midway is a reproduction of the far-famed Street of All Nations of 1893, with many improvements upon the original.

The electric lighting of the buildings, grounds, and lagoon at night makes a scene of enchanting beauty, alone worth traveling a thousand miles to see.

The means of communication between

the city and the grounds are ample, and the distance to be traversed is short.

The ways of reaching Omaha are innumerable, but chief among them is the direct Chicago and Omaha short line of the Chicago, Malwaukee & St. Paul Railway, with its electric-lighted, vestibuled trains, leaving Chicago every night at 6.15 P.M., and arriving Omaha at 8.20 the next morning. Dining-car service *en route*.

Excursion-tickets are on sale at every coupon ticket office in the United States over the Chicago, Milwaukee & St. Paul Railway through Northern Illinois and

PURE AND RELIABLE ANIMAL VACCINE LYMPH.
Fresh Daily. Send for Illustrated Pamphlet.
10 IVORY POINTS, Double Charged, \$1.00. 10 CAPIL-
LARY TUBES, \$1.00. Orders by Mail or Tele-
graph Promptly Dispatched.

VACCINE**VIRUS.**

NEW ENGLAND VACCINE CO., Chelsea Station, BOSTON, MASS.

WM. C. CUTLER, M.D.

CHAS. N. CUTLER, M.D.

AS AN ANODYNE AND HYPNOTIC

...KRYOFINE...

(Methoxacet-p-phenetidin)

IS POWERFUL AND WITHOUT ILL EFFECTS.

DR. GEORGE F. BUTLER, Professor of Materia Medica and Clinical Medicine, College of Physicians and Surgeons, Chicago; Attending Physician Cook County Hospital, etc., in the course of an article in the Chicago Clinic, says:

"With the exception of morphine, no drug possesses so positive, prompt and efficient an analgesic property as Kryofine; indeed there are painful disorders—such as migraine, and particularly the pains of locomotor ataxia and certain spinal diseases—where Kryofine seems nearly as efficient as morphine, and attended with less unpleasant sequelæ."

In the services of Drs. J. Rudisch, A. Meyer and A. G. Gerster, Mt. Sinai Hospital, New York, it was found that:

"As a hypnotic in insomnia, unaccompanied by severe pain, it has proved of undoubted value."

"In a case of acute suppurative arthritis, general sepsis and pericarditis, the drug did not affect the pulse. When the child was restless at night, 2½ grains of Kryofine produced a quiet sleep, lasting several hours."—DRS. HAAS and MORRISON, New York Med. Journal.

Dose:—4-7½ grs., tablets or crystalline powder.

Literature and Samples sent on request.

C. BISCHOFF & CO., NEW YORK.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

Central Iowa, as well as at 95 Adams St., and at the Union Passenger Station, Canal and Adams streets, Chicago.

The Action of Ze-an in Renal and**Urinary Disorders.**—The importance

of the eliminative function of the kidneys cannot be overestimated. It is the complement to that of assimilation. The food may be rich in nutritive elements, but if the process of digestion be not thoroughly performed deleterious substances will accumulate in the blood. It is highly essential, therefore, that they should as speedily as possible be removed from the blood and system. The work of this elimination falls largely upon the kidneys. These delicate glandular organs are, in a state of health, amply competent to the task which nature has laid upon them and keep the blood free from impurities. When,

through any cause or conjunction of causes waste-products are increased, the kidneys are stimulated to unusual or excessive efforts. This excess of functional activity reacts, in time, upon the structure of the organs. The cells of the kidneys undergo pathological alterations, and their efficiency is crippled or altogether destroyed. A vast series of morbid conditions is the consequence of this incompetency of the kidneys.

Many of the waste products which circulate in the blood are insoluble in watery fluids, and consequently as they pass out of the body in the urine they irritate the urinary organs. If they are deposited in the kidney they form calcareous concretions. If they gain the bladder before deposit they give rise to stone in the bladder. The passage of smaller particles as sand or gravel is a frequent cause of inflammation of the kidney or bladder.

ELIXIR MALTOPEPSINE

(TILDEN'S)

DIGESTS ALL KINDS OF FOOD.

Is especially useful in **CHOLERA INFANTUM** and all forms of **SUMMER COMPLAINT WITH CHILDREN**. Promotes digestion by aiding and perfecting the process. Gives immediate relief in **DYSPEPSIA**, and aids in curing it. Send for samples and literature.

RESPIRAZONE

(TILDEN'S)

Indispensable in the treatment of **ACUTE** and **CHRONIC ASTHMA**, **HAY FEVER**, and **CROUP**. Physicians everywhere are prescribing this preparation with unvarying results. It is not a new remedy, but a correct combination of old remedies, and the profession can prescribe it feeling confident of results. Doctor, if you have not used **RESPIRAZONE** (TILDEN'S) send at once for samples and literature.

MANUFACTURED ONLY BY

THE TILDEN COMPANY,

MANUFACTURING PHARMACISTS,

NEW LEBANON, N. Y.

ST. LOUIS, MO.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

Other organic compounds which are retained in the circulation when the kidneys are disabled lay the foundation of disease of the heart and blood-vessels. Another class of waste-products act as poisons to the cells of the nervous system.

In treating renal diseases, therefore, we must, in the first place, regulate the diet in such a manner that formation of noxious waste-products shall be reduced to a minimum. In the next place we must take measures to convert insoluble into soluble substances. If we decrease the production of insoluble bodies and if we sweep rapidly out of the system those whose generation we cannot prevent, we have evidently done much toward bringing about recovery. The third element of the problem is to soothe the irritability, congestion, or inflammation of the kidney or bladder.

A remedy which possesses many of the properties desirable in treating the disorders of which we have been speaking is corn-silk. It is a mild stimulant to the kidneys, a solvent for uric acid and phosphatic calculi and it also exerts a certain analgesic effect upon the entire urinary tract. The official preparations of corn-silk, however, labor under the disadvantage of containing an insufficient proportion of the active principle. For this reason ze-an is offered to the medical profession as an acceptable form in which to prescribe corn-silk. Ze-an is a pure and concentrated fluid extract of corn-silk. Its strength renders it efficient in drachm-doses.

By promoting the solution of gravel ze-an prevents the occurrence of those painful and exhausting attacks of renal colic which are caused by the passage of insoluble and, consequently, irritating particles along the ureter and into the

White Rock Ozonate Lithia Water

PLEASES PARCHED PALATES PERFECTLY.



*People drink it because
they like it; hence the uniform
success reported by thousands of
physicians who prescribe it.*

Bottled Only by

WHITE ROCK MINERAL SPRINGS CO., Waukesha, Wis.

In writing to advertisers mention Monthly Cyclopædia

Notes and Comments (continued).

bladder. Continuing the same action the formation of stone in the bladder is likewise prevented.

The irritant action of gravel and stone upon the kidney and bladder gives rise to pain and inflammation in those organs and, in the kidney, the trouble not infrequently proceeds to suppuration. In renal congestion, nephritis, and pyelitis the valuable combination of properties present in corn-silk renders ze-an a reliable remedy. This preparation, for similar reasons, is equally serviceable in chronic cystitis whether due or not to the presence of a foreign body. Prostatitis and urethritis are also relieved by the use of the same remedy. In fact, there is strong evidence to the efficacy of ze-an in acute gonorrhœa. This preparation has been administered with success in cases of nocturnal incontinence of urine.

The diuretic properties of ze-an have been both experimentally and clinically proved. The remedy is serviceable in dropsy dependent upon either cardiac or renal disease. By removing the transudation ze-an acts indirectly as a tonic to the heart.

Karozza: Ethics of Marriage. By Alice B. Stockham, M.D., author of "Tokology." Published by Alice B. Stockham & Co., 277 Madison Street, Chicago. Special price to physicians, \$1.00, prepaid.

This work was written from years of professional experience to meet a demand among inquiring women on subjects deeply concerning the physical life of the wife and mother. It elucidates the theory of conjugal life on a peculiar subject, and states plain facts in plain language, and will be interesting reading to many who hold to the Malthusian

doctrine, and the legitimate limitation of families. Sent, postage paid, on the receipt of price to the publishers.

American Tapestry Decorative Company.

—The firm of John F. Douthitt, 286 Fifth Avenue, New York, carries on a large decorative business in many directions. Tapestry materials, goblin-printed burlaps, and goblin art drapery are among the firm's products. The company will send books and illustrations of their goods for the benefit of those interested in artistic decoration. In connection with the business a school is also maintained in which tapestry-painting lessons are given. Complete written instructions will be sent by mail.

Kryofine.—Kryofine is an efficient agent in reducing fever and alleviating pain. Not only does it lower a febrile temperature, but, if given prior to the usual time of a rise of the curve, it prevents the expected ascent. In typhoid fever and influenza it has been used with very satisfactory results. In rheumatic fever it is doubly valuable on account of its effect both on the temperature and pain. Kryofine is a valuable remedy in the various forms of neuralgia. It has been employed with advantage also in polyneuritis.



**DR. BECKER'S
COMPOUND
DIGEST.**

(*Digestinla Composita. Dr. Becker.*)

is prepared according to the formula of Dr. Becker, from principles obtained by his processes, from the digestive organs of clean animals and fowls.

It assists impaired and overworked digestive organs, allowing them to rest without lessening nutrition.

It is easily administered, and, being free from chemicals, does not interfere with the action of other medicines.

Sold only in bottles containing 550 grains powder form. or 100 five-grain Tablets, and may be ordered through any druggist. Samples sent by mail to physicians on application.

DR. BECKER COMPOUND DIGEST CO.,

107 Dearborn Street, CHICAGO, ILL.

In writing to advertisers mention Monthly Cyclopædia

THE SANITARIUM, BATTLE CREEK, MICH.

INCORPORATED 1867.



THE largest, most thoroughly equipped and one of the most favorably located in the United States. It is under strictly regular management. Eight physicians, well-trained and of large experience. A quiet home like plan with "trains and courses," "rest-cure," "massage," "faradization," "galvanization," "static electricity," "Swedish movements," "dieting," "laths," "physical training," and all that pertains to modern rational medical treatment can be had in perfection at reasonable prices. Special attention given to the treatment of chronic disorders of the stomach, and diseases peculiar to women. A special hospital building (too beds) for surgical cases with finest hospital facilities and appliances.

Large fan for winter and summer ventilation. Absolutely devoid of usual hospital odors. Delightful surroundings. Lakeside resort. Pleasure grounds, steamers, sailboats, etc. Trained nurses of either sex furnished at reasonable rates. **I. H. KELLOGG, M. D., Superintendent, Battle Creek, Mich.**

J. H. KELLOGG, M. D., Supt., Battle Creek, Mich.

NUT-GLUTEN BISCUIT: Made from the purest wheat gluten and nut meal. The only food on the market to-day that diabetics may eat of freely, without danger. Prescribed and endorsed by the physicians of the Battle Creek Sanitarium. Samples free to physicians when card enclosed with the request. Others may have them for two 3c stamps.

BATTLE CREEK SANITARIUM HEALTH FOOD CO., Battle Creek, Mich.



In writing to advertisers mention Monthly Cyclopædia

Artistic Home Decorations.

We can show you effects **NEVER** before thought of, and at moderate prices, too.

Why have your house decorated and painted by inferior workmen, when you can have it done by skilled workmen—by Artists—for the same price?

If you intend Decorating, if only one room, call or send to see what we are doing, and for whom.

TAPESTRY PAINTINGS.

2000 tapestry paintings to choose from. 38 artists employed, including gold medalists of the Paris Salon. Send 25 cents for compendium of 140 studies. When in the city be sure to call and see our exhibition.

WALL PAPERS.

New styles, designed by gold-medal artists. From 10 cents per roll up. Send 50 cents to prepay expressage on large sample books and drapery. Will include drapery samples in package. See our Antique Metallic, French, Pressed Silks, and Lida effects in special colors to match all kinds of wood-work, carpets, and draperies. Have 500 different wall-hangings with draperies specially made at our Broomhead Mills, Paterson, N. J., to match.

DECORATIONS.

Color-schemes—designs and estimates submitted free. Artists sent to all parts of the world to do every sort of decorating and painting. We are educating the country in color-harmony. Relief, stained glass, wall-paper, carpets, furniture, draperies, etc. Pupils taught.

DECORATIVE ADVICE.

Upon receipt of \$1 Mr. Douthitt will answer any question on interior decorations—color-harmony and harmony of form, harmony of wall-coverings, carpets, curtains, tiles, furniture, gas-fixtures, etc.

MANUAL OF ART DECORATIONS.

The art-book of the century. 200 royal quarto pages, filled with full-page colored illustrations of modern home interiors and tapestry studies. Price \$2.

If you want to be up in decoration send \$2 for this book; worth \$50.

SCHOOL.

Six 3-hour tapestry painting lessons, in studio, \$5. Complete written instructions by mail, \$1. Tapestry paintings rented; full-size drawings, paints, brushes, etc., supplied. Nowhere, Paris not excepted, are such advantages offered pupils. New catalogue of 125 studies, 25 cents.

Send \$1 for complete instructions in tapestry painting and compendium on 140 studies.

TAPESTRY MATERIALS.

We manufacture tapestry materials superior to foreign goods and half the price. Book of samples, 10 cents.

Send \$1.50 for 2 yards No. 6, 50-inch goods, just for a trial order; worth \$3. All kinds of draperies to match all sorts of wall papers, from 10 cents per yard up. **THIS IS OUR GREAT SPECIALTY.**

GOBLIN-PRINTED BURLAPS.

Over 100 new styles for wall-coverings, at 25 cents per yard, 35 inches wide, thus costing the same as wall-paper at \$1 per roll. 240 kinds of Japanese lida leather papers at \$2 per roll.

GOBLIN ART DRAPERY.

Grecian, Russian, Venetian, Brazilian, Roman, Rococo, Dresden, Festoon, College Stripe, Marie Antoinette, Indian, Calcutta, Bombay, Delft, Soudan.

In order that we may introduce this line of New Art Goods, we will send one yard each of 50 different kinds of our most choice Patterns for \$7.50.

JOHN F. DOUTHITT,

American Tapestry Decorative Co.,

286 Fifth Avenue, New York.

Near 30th Street.

In writing to advertisers mention Monthly Cyclopaedia

1898.

WE HAVE JUST ISSUED

— A New 32-Page —

DESCRIPTIVE CATALOGUE



OF ALL OUR PUBLICATIONS,

Which will be cheerfully mailed
to any address promptly
on request.

THE F. A. DAVIS COMPANY,

MEDICAL PUBLISHERS.

PHILADELPHIA: 1914 and 1916 Cherry Street.

NEW YORK: 117 W. Forty-Second Street

CHICAGO: 9 Lakeside Building, 214-220 S. Clark St.

CLIMATE CURE

the SALT RIVER VALLEY of Arizona and the various HEALTH RESORTS in NEW MEXICO

ARE UNRIVALED FOR THE RELIEF OF CHRONIC LUNG AND THROAT DISEASES. HERE ARE FOUND PURE DRY AIR, EQUABLE TEMPERATURE AND CONSTANT SUNSHINE. THE ITEMS OF ALTITUDE, TEMPERATURE, HUMIDITY, HOT SPRINGS, SANATORIA, COST OF LIVING, MEDICAL ATTENDANCE, SOCIAL ADVANTAGES, ETC., ARE CONCISELY TREATED IN DESCRIPTIVE PAMPHLETS ISSUED BY THE SANTA FE ROUTE. PHYSICIANS ARE RESPECTFULLY ASKED TO PLACE THIS LITERATURE IN THE HANDS OF INVALIDS.

ADDRESS W.J. BLACK,
G.P.A., A.T. & S.F. RY.,
TOPEKA, KANSAS.

OR C.A. HIGGINS,
A.G.P.A., CHICAGO.

NOW READY.

A Very Important Contribution to Laryngological Science.

Autoscopy of the Larynx and the Trachea.

(DIRECT EXAMINATION WITHOUT MIRROR.)

By **ALFRED KIRSTEIN, M.D.**, Berlin.

Authorized Translation (Altered, Enlarged, and Revised by the Author) by

MAX THORNER, A.M., M.D., Cincinnati, O.

Professor of Clinical Laryngology and Otology, Cincinnati College of Medicine and Surgery; Laryngologist and Aurist, Cincinnati Hospital, etc.

About 75 Pages, with Twelve Illustrations. Crown Octavo, bound in Extra Cloth, price, 75 cents net.

THE F. A. DAVIS COMPANY, Publishers.

PHILADELPHIA: 1914 and 1916 Cherry St.

NEW YORK: 117 W. Forty-Second St.

CHICAGO: 9 Lakeside Building, 214-220 S. Clark St.

In writing to advertisers mention Monthly Cyclopædia

PEPPER WHISKY

ESTABLISHED 1780

THE GENUINE HAS THIS COUPON



PULL OFF ONLY
When Bottle is
"PULLED"
PATENTED

EXTRA CAUTION
Anyone refilling this bottle without destroying the label is criminally liable and on detection will be prosecuted to the full extent of the law

DISTILLERY BOTTLING
OLD PEPPER WHISKY
1780

THE OLDEST & BEST BRAND OF WHISKY MADE IN KENTUCKY

PEPPER
DISTILLERY
HAND MADE
SOUR MASH
ESTABLISHED 1780
JAS. E. PEPPER & CO.
DISTILLERS,
LEXINGTON, KY.

TRADE MARK

Jas. E. Pepper & Co.
LEXINGTON, KY.

READ COUPON ON BACK OF THIS LABEL

BUY
GENUINE
DISTILLERY
BOTTLING
ONLY

FROM
HONEST DEALERS
OR THE
DISTILLERY DIRECT

AND THUS
AVOID IMITATIONS
AND REFILLED
BOTTLES

Genuine distillery bottling of Old Pepper Whisky, distilled only by Jas. E. Pepper & Co., Lexington, Ky., under the same formula for more than 100 years, is guaranteed absolutely the purest and best in the world. Sample case, \$15.00, sent on trial, which, if not perfectly satisfactory, can be returned and money will be refunded.

COPYRIGHT 1897 BY
J. E. PEPPER & CO.
ST. LOUIS, MO.
CINCINNATI & NEW YORK

Read and save coupons on genuine Distillery Bottling of Old Pepper Whisky and see who gets the \$5500 in 1898, in addition to \$1.00 per dozen.

Been giving satisfaction for 118 years---need we say more ?

In writing to advertisers mention Monthly Cyclopædia



Dietz Driving Lamp

IT is the only perfect one.
 IT throws all the light straight ahead
 from 200 to 300 feet.
 IT looks like a locomotive headlight.
 IT gives a clear white light.
 IT burns kerosene (Coal Oil)

It will not blow nor jar out



SPECIAL OFFER. CUT THIS ADVERTISEMENT OUT and send it to us and we will send book describing our lamp, and will agree to send you one single lamp or a pair at our wholesale price (very much less than the retail price).

R. E. DIETZ COMPANY, 60 Laight St., New York.
 ESTABLISHED 1840.

WAMPOLE'S LIQUID WHEAT.

Each teaspoonful represents **one ounce** of all parts of the **whole wheat grain**, partly pre-digested and concentrated *in vacuo*; has sufficient **diastase** to digest five times its weight of dry starch, and contains one-eighth grain U. S. P. Pepsin.

Liquid Wheat contains over **seven per cent.** of the **Nutrient Organic Hypophosphites**, obtained solely from wheat, principally from the germ and cortical portions. Contains all the elements of nutrition, and is an efficient aid to the digestion of both **starchy** and **nitrogenous** foods. Valuable as a supplementary diet for growing children, insuring sufficient **tooth nutrition**, allaying **nervous irritability**, and inducing **plumpness**.

SAMPLES AND LITERATURE CHEERFULLY FURNISHED UPON APPLICATION.

Prepared by

HENRY K. WAMPOLE & CO.,

Manufacturing Pharmacists, Philadelphia, Pa.

"DO NOT STAMMER."

Philadelphia, Pa., June 18, 1898.



EDWIN S. JOHNSTON,
Principal and Founder

DEAR MR. JOHNSTON:

I count it a privilege to bear testimony to the efficacy of the method employed by you for the cure of speech defects, also to the very high character of your most successful Institute.

It is more than six years since, under your tuition I was cured of a distressing impediment of over fifteen years' standing. I have been almost continuously engaged in teaching since leaving your school, and I have, during the past three years or more, been in the work of the Christian ministry in this city. There is no doubt about the permanency of my cure. I am acquainted with a number of persons whose permanent cures are of longer standing than my own.

More than this, I owe much to the mental discipline and to the moral and spiritual help received while under your tuition. The deep personal interest you have always shown in the welfare of your pupils is one of the commendable features of your school. No parent need hesitate about sending his child to you.

I shall never be able to fully express my gratitude for what has, in the providence of God and through your instrumentality, been accomplished in me. Ever most gratefully yours,

CASPAR C. GARRIGUES,
Ass't Pastor First Church of Christ,
2716 "A" St., Phila.

Send for 60-page book to the PHILADELPHIA INSTITUTE, 1033 Spring Garden St., Philadelphia, Pa. Established 1884. **EDWIN S. JOHNSTON**, Principal and Founder (who cured himself after stammering 40 years).

JUST READY. New, Revised Edition of an exceedingly useful ... book ...

A Primer of Psychology and Mental Disease

FOR USE IN TRAINING-SCHOOLS FOR ATTENDANTS AND NURSES
AND IN MEDICAL CLASSES.

By **C. B. BURR, M.D.,**

Medical Director of Oak Grove Hospital for Nervous and Mental Diseases, Flint, Mich. Formerly Medical Superintendent of the Eastern Michigan Asylum. Member of the American Medico-Psychological Association, etc.

Second Edition. Thoroughly Revised and Illustrated. 116 pages, 5½ x 8 inches. Neatly Bound in Extra Cloth. Price, \$1.00, Net, POST-PAID.

THE F. A. DAVIS COMPANY, PUBLISHERS, 1914 AND 1916 CHERRY STREET, PHILADELPHIA.

In writing to advertisers mention Monthly Cyclopædia

WILL SOON BE PUBLISHED.

INJURIES TO THE EYE
IN THEIR
MEDICO-LEGAL ASPECT.

By S. BAUDRY, M.D.,

PROFESSOR IN THE FACULTY OF MEDICINE, UNIVERSITY OF LILLE, FRANCE, ETC.

TRANSLATED WITH THE PERMISSION OF THE AUTHOR AND EDITED BY

CHARLES A. OLIVER, A.M., M.D.,

ONE OF THE ATTENDING SURGEONS TO THE WILLS EYE HOSPITAL, PHILA.; ONE OF THE OPHTHALMIC SURGEONS TO THE PHILADELPHIA HOSPITAL; MEMBER OF THE AMERICAN AND FRENCH OPHTHALMOLOGICAL SOCIETIES, ETC.

Crown Octavo. About 200 pages. Bound in Extra Cloth. Price, \$1.25, net.

This work, which is undoubtedly the most able and important modern contribution in any language on the subject of Injuries to the Eye in Their Medico-Legal Relations, and only quite recently issued in France, is now for the first time adapted to the American and English profession. It treats the subject in such a thorough and practical way that we feel sure it must be cordially welcomed by the profession of both England and America.

THE F. A. DAVIS COMPANY Medical Publishers.

PHILADELPHIA: 1914 and 1916 Cherry Street.

NEW YORK: 117 West Forty-Second Street.

CHICAGO: 9 Lakeside Building, 214-220 South Clark Street.

A MOST INTERESTING NEW BOOK.

AN EPITOME OF THE
HISTORY OF MEDICINE

From the Earliest Times to the Present.

By ROSWELL PARK, A.M., M.D.,

Professor of Surgery and Clinical Surgery in the Medical Department of the University of Buffalo, Buffalo, New York, etc., etc.

Handsomely Illustrated with Half-tone Reproductions of Portraits of the Eminent Physicians and Surgeons of all time and other Engravings.

Royal Octavo. 350 Pages. Extra Cloth. \$2.00 net. Now Ready.

THE F. A. DAVIS CO., Publishers, 1914-16 Cherry St., Philadelphia.

In writing to advertisers mention Monthly Cyclopædia

Salesmen Wanted.

Years ago the country merchant went to the city once or twice a year to buy his goods. Now a traveler calls on him every week, and he buys what he wants from the latest samples.

The traveler has come to stay.

Years ago the country doctor received catalogues and ordered therefrom, when money was so plentiful that he did not know what else to do with it. Consequently he did not order very much, and good libraries were few and far between.

Following the example of the dry-goods man, the medical publisher began sending out travelers about twenty-five years ago. Is the universally admitted fact, that medicine has made greater progress in that time than during a century before, a coincidence only?

However this may be, the fact is that the progressive country doctor, a hundred miles from any city, now keeps better informed than the city doctor did then.

He has become accustomed to having the latest ideas in medicine presented to him by the wide-awake traveler. He welcomes him as a source of reliable information on the best publications.

While he does not always buy, he has, like the dry-goods man, become accustomed to consider this method of buying the most satisfactory one, because he can examine his purchase thoroughly before he pays his money for it.

In this is the traveler of great use to the physician.

We claim to have the largest force of *expert* travelers in the business. We want a few more. Physicians who would prefer to hustle days rather than nights, can hear about it by addressing, with particulars,

THE SUBSCRIPTION DEPARTMENT,

THE F. A. DAVIS COMPANY,

1914-16 Cherry St., Philadelphia, Pa.

In writing to advertisers mention Monthly Cyclopædia

ATTENTION! INVALIDS AND PLEASURE-SEEKERS.
There is but ONE place TO FIND
HEALTH AND PLEASURE,
-- COLORADO --

THE DENVER & RIO
GRANDE RAILROAD.

"Scenic Line of
the World."



The only line of rail-
road reaching all Col-
orado's many beautiful
resorts.

THE LAND OF CLOUDLESS DAYS AND BALMY NIGHTS.

For Illustrated Descriptive Matter and Information as to Rates and
Routes, call on or address,

S. K. HOOPER, G. P. AND T. A., DENVER, COLO.

12th YEAR.

R. L. POLK & CO.'S
Medical and Surgical Register
OF THE UNITED STATES AND CANADA.

This is positively the only national Medical Directory published.

Embraces names of over 112,000 Physicians, with college of graduation, list of Colleges, Societies, Boards of Health, Journals, Mineral Springs, Hospitals, Sanitariums, Asylums, and other Medical Institutions; also Medical Laws of each State.

If you have anything to say to the Physicians in the United States and Canada there is no better or more economical way of saying it to them than by an advertisement in the Register; or if you wish to send them your circulars, the Register contains the list in convenient form for addressing.

R. L. POLK & CO.,

New York.

Chicago.

Baltimore.

Detroit.

In writing to advertisers mention Monthly Cyclopædia

JUST PUBLISHED.

EYE-STRAIN IN HEALTH AND DISEASE,

WITH SPECIAL REFERENCE TO THE

AMELIORATION OR CURE OF CHRONIC NERVOUS DERANGEMENTS
WITHOUT THE AID OF DRUGS.

A Manual for the General Practitioner in Medicine to Aid in Detecting the Presence
and Appreciating the Importance of Eye-strain in
Health and Disease.

By AMBROSE L. RANNEY, A.M., M.D.,

AUTHOR OF "LECTURES ON NERVOUS DISEASES," "THE APPLIED ANATOMY OF THE NERVOUS SYSTEM," "A
TREATISE ON SURGICAL DIAGNOSIS," "PRACTICAL MEDICAL ANATOMY," ETC. LATE PROFESSOR OF
ANATOMY OF THE NERVOUS SYSTEM IN THE NEW YORK POST-GRADUATE MEDICAL SCHOOL
AND HOSPITAL; LATE PROFESSOR OF NERVOUS DISEASES IN THE MEDICAL
DEPARTMENT OF THE UNIVERSITY OF VERMONT, ETC., ETC.

THIS treatise is one of so much importance to the general practitioner that its great practical value can hardly be overestimated. For many years Dr. Ranney has devoted the most painstaking and unprejudiced study and investigation to an almost endless variety of cases. The results are crystallized and given unreservedly to the medical profession in this handy volume.

The various chapters of the work are replete with illustrative cases. They discuss the results of Eye-treatment of Headache, Neuralgias, Sleeplessness, St. Vitus's Dance, Chronic Digestive Diseases, Epilepsy, Nervous Prostration, and Insanity.

The steps and instruments employed in testing by the latest methods the eyes and eye-muscles are described with sufficient detail to meet the requirements of the general practitioner in medicine, as well as the oculist, in making a diagnosis.

The indications and contra-indications for operative procedures upon the eye-muscles, and the method of performing graduated tenotomies and advancement operations are furthermore discussed in a separate chapter.

Fully Illustrated. In One Royal Octavo Volume. 325 Pages. Handsomely Bound in Extra Cloth, \$2.00 net.

THE F. A. DAVIS CO., PUBLISHERS,
1914 AND 1916 CHERRY STREET, PHILADELPHIA, PA.

In writing to advertisers mention Monthly Cyclopædia

We are
Determined *that the*

\$3.50 FOR \$1.75.

FALL OF 1898 SHALL BE A MEMORABLE ONE

IN THE HISTORY OF

The Medical Bulletin

By a Great Increase in its Already Large Subscription List. To do this we make the following hitherto

UNEQUALED PREMIUM OFFER:

Wishing to be fair to all, and not deeming it just (though our principal aim is new subscribers) to deny our old friends equal advantages, but, on the other hand, considering them fully entitled to the best of all, we simply say that, so long as the present supply of this valuable book exists, it will be furnished to all who send in their subscriptions for the "**Bulletin**," whether they be old or new subscribers. The book we offer is

OBSTETRIC SURGERY,

By ECBERT H. GRANDIN and GEORGE W. JARMAN, M.D.

Royal Octavo, 250 pages, 85 engravings in the text, and 15 handsome full-page Photographic plates, Bound in extra Cloth, Published and sold at \$2.50 net, which will be furnished, shipping charges paid by the publishers, including a full year's subscription to the "**Medical Bulletin**" (regular subscription-price, \$1.00 per year),

For \$1.75, Cash with Order.

This most excellent, practical volume was recently issued as a part of the same authors' complete work, entitled "Practical Obstetrics," and as soon as our present limited supply is exhausted no more can be obtained.

The "Medical Bulletin" points with pride to its great record in 1897. A brief review of its work during that year may be interesting. During the year 1897 the "Bulletin" printed ninety (90) original articles by distinguished authorities, embracing the whole broad domain of medicine and surgery. Its illustrated pathological articles by the celebrated Dr. Isaac Ott have attracted world-wide attention. It has published no less than 375 square inches of illustrations: more than equal to that contained in the average \$4.00 book. It has published no less than 250 approved and practical formulæ, all having passed under the critical eye of its distinguished editor, Dr. John V. Shoe-

In writing to advertisers mention Monthly Cyclopædia

maker, whose great work on "Materia Medica and Therapeutics," now in its Fourth Edition, and used in a hundred colleges, is known and read all over the English-speaking world. This portion of the "**Bulletin**" alone, published in book form, would comprise a "Formulary" far in advance of many on the market, whose price is double that which has been charged the fortunate subscribers to the "Bulletin." It has kept its readers fully informed as to college and society happenings; has given hundreds of timely hints, which have been of great service to its readers. Has published a series of articles relating to that great "American Riviera," the "West Coast of Florida," that has been of incalculable service to its readers in aiding them to intelligently advise their consumptive and rheumatic patients upon economic as well as hygienic topics.

It has given its readers no less than 350,000 words of matter equal to the usual contents of a 700-page book, selling at an average price of \$4.00, as, for instance, that excellent book: Gray, on **Nervous Diseases**.

But the "**Bulletin**" purposes to do even more during the coming year, and in addition give its friends an opportunity seldom equaled in journalism.

Old subscribers can have their subscriptions advanced in availing themselves of this splendid offer. New subscribers can have their subscriptions begin with the first of January, 1899, or sooner if they prefer. **Whether you are an old or a new subscriber all you have to do is to send us cash to the amount of \$1.75 and promptly receive a copy of Drs. Grandin and Jarman's "Obstetric Surgery" and the "Bulletin" will be sent you regularly for one year.**

It is important to act promptly; therefore we urge your sending your subscription immediately with P. O. or Express Money-Order, or check, to

THE F. A. DAVIS COMPANY, Medical Publishers,

1914 and 1916 Cherry Street, Philadelphia, Pa.

SPECIAL ORDER BLANK.

FILL THIS OUT AND MAIL IT, WITH \$1.75, TO THE PUBLISHERS,

THE F. A. DAVIS COMPANY,

1914 and 1916 Cherry Street, Philadelphia.

Gentlemen: Enclosed find One Dollar and Seventy-five Cents (\$1.75), for which please send me, all carriage charges prepaid, a copy of "OBSTETRIC SURGERY," by DRs. GRANDIN and JARMAN, Complete, Bound in Extra Cloth (Regular price, \$2.50 net), and "THE MEDICAL BULLETIN" (regular subscription-price, \$1.00 per year) for one year, as per your Premium Offer for the months of August and September, 1898.

Name _____

P. O. Address _____

County _____

State _____

P. S.—Upon the reverse side of this announcement will be found a specimen page of Grandin and Jarman's "Obstetric Surgery."

In writing to advertisers mention Monthly Cyclopædia

Specimen page from Grandin and Jarman's "Obstetric Surgery."

VERSION.

111

As the cord comes down it is to be drawn upon from the *placental* side, and if it is over one of the legs it must be released (Fig. 53) and placed in the most favorable position as regards pressure. In rare instances it will be impossible to draw the cord down without making undue traction. If such should prove to be the case, it should be secured by means of two artery-clamps and cut. Of course, if this is done, it will be necessary to hasten the delivery as much as possible.

When the scapulæ appear the arms must be liberated before extraction is continued. Under favorable circumstances,—that is, if the assistant has kept up intelligent pressure on the fundus, or if the cervix was fully dilated previous to the



Fig. 53.—Method of Releasing the Cord.

version, or if the operator has not made traction in too rapid a manner,—the arms will be folded on the chest and their extraction will be easy.

Even in the hands of the best operators and with the best assistants the arms sometimes become unavoidably extended. Although their extraction must be accomplished in as rapid a manner as possible, there is no need of breaking the arm if care is taken.

The arm which is to the rear is usually more easily liberated. To do this the operator seizes the legs with one hand and carries the child's body well upward. This will cause the posterior shoulder to be more readily reached, and will permit more room for the manipulations necessary. Two fingers of

IMPORTANT ANNOUNCEMENT

*Concerning DR. WINFIELD S. HALL'S Translation of the Latest German Edition
of Edinger's "Structure of the Central Nervous System."*

IN PRESS. WILL BE READY SOON.

THE ANATOMY OF THE Central Nervous System of Man and of Vertebrates in General.

By Prof. LUDWIG EDINGER, M.D.,
Frankfort-on-the-Main, Germany.

Translated from the FIFTH GERMAN EDITION,

By WINFIELD S. HALL, Ph.D., M.D.,
Professor of Physiology in the Northwestern University Medical School, Chicago.

ASSISTED BY

PHILO LEON HOLLAND, M.D., Instructor in Clinical Neurology, and
EDWARD P. CARLTON, B.S., Demonstrator of Histological Neurology in the Northwestern
University Medical School, Chicago.

Professors and Teachers of Anatomy of the Nervous System, Histology, etc., in Medical Colleges throughout the English-speaking world will be gratified to learn that the above book will be ready in time for the college sessions of '98 and '99.

This translation is a thoroughly complete one from the latest German edition, admirably executed, and done into clear, correct, and readable English with the authority and indorsement of the famous author. It contains upward of 250 engravings in the text and comprises over 400 Royal Octavo pages. It will be handsomely printed and bound in extra cloth, and sold at the exceedingly moderate price of \$2 50 net.

That this is one of the most important, interesting, and useful books which can be placed in the hands of a medical student during his attendance at college will be quickly recognized by every Teacher of Anatomy, Histology, and Neurology.

Further information promptly furnished by the publishers.

THE F. A. DAVIS COMPANY, Publishers.

PHILADELPHIA: 1914 and 1916 Cherry St.
NEW YORK: 117 W. Forty-Second St.
CHICAGO: 9 Lakeside Building, 214-220 S. Clark St.

In writing to advertisers mention Monthly Cyclopædia

SYR. HYPOPHOS. CO., FELLOWS

Contains the Essential Elements of the Animal Organization—Potash and Lime;

The Oxidising Agents—Iron and Manganese;

The Tonics—Quinine and Strychnine;

And the Vitalizing Constituent—Phosphorus; the whole combined in the form of a Syrup with a Slightly Alkaline Reaction.

It Differs in its Effects from all Analogous Preparations; and it possesses the important properties of being pleasant to the taste, easily borne by the stomach, and harmless under prolonged use.

It has Gained a Wide Reputation, particularly in the treatment of Pulmonary Tuberculosis, Chronic Bronchitis, and other affections of the respiratory organs. It has also been employed with much success in various nervous and debilitating diseases.

Its Curative Power is largely attributable to its stimulant, tonic, and nutritive properties, by means of which the energy of the system is recruited.

Its Action is Prompt; it stimulates the appetite and the digestion, it promotes assimilation, and it enters directly into the circulation with the food products.

The prescribed dose produces a feeling of buoyancy, and removes depression and melancholy; *hence the preparation is of great value in the treatment of mental and nervous affections.* From the fact, also, that it exerts a double tonic influence, and induces a healthy flow of the secretions, its use is indicated in a wide range of diseases.

Medical Letters may be addressed to **Mr. FELLOWS, 48 Vesey Street, New York.**



Open All the Year. 2,500 FEET ELEVATION.
ON CHESAPEAKE & OHIO RAILWAY.

THE NEW HOMESTEAD,

a grand new hotel completed last year, equipped and furnished throughout in accordance with the latest designs and most modern improvements. The Virginia, the Annex, and ten new cottages clustered near the Homestead. A handsome new Casino adds greatly to indoor enjoyment.

*Golf Links and Club House, Tennis Courts, Billiards, Pool,
Riding School and Fine Livery.*

MAGNIFICENT MOUNTAIN SURROUNDINGS.

FINEST BATH-HOUSE IN AMERICA.
MOST CURATIVE BATHS KNOWN

for rheumatism, gout, and nervous troubles. All baths from flowing streams of natural hot water.

Located nearly midway between Washington, Old Point Comfort, and Cincinnati or Louisville. Hot Springs is an exceptionally convenient and delightful stopping-place for west-bound travelers.

Special Pullman Compartment Car leaves Pennsylvania Station, New York, 4 50 p.m., Tuesday, Thursday, and Saturday, and arrives at Hot Springs via Washington, 7 25 a.m. Dining car.

Excursion tickets, baggage checks, and Pullman Reservations at Chesapeake & Ohio Railway Offices, 362 and 1323, Broadway, New York, also offices Pennsylvania Railroad and connecting lines.

For illustrated pamphlets and further particulars, address W. H. GREGOR, General Agent Passenger Department Chesapeake and Ohio Railway, Washington, D. C., or

FRED STERRY, Manager, Hot Springs, Bath Co., Va.

In writing to advertisers mention Monthly Cyclopædia

Oak Grove Hospital


For Nervous and Mental Diseases.



Equipment for hydrotherapeutic and electrical treatment complete and modern—Static, Galvanic, and Faradic Apparatus, Electric Bath, Turkish and Russian Baths, and Massage. Noyes Amusement Hall in connection with the Hospital, containing gymnasium, billiard-room, bowling-alley, hall for assemblies, sitting-room, lounging-room, and library, furnishes ideal facilities for in-door sports and diversion. Carriages for the daily use of patients free. For terms, address

Grounds comprise sixty acres of stately oaks and are picturesque and secluded. Buildings roomy, homelike, free from institutional features, and erected with especial reference to the care of nervous and mental invalids.

Interiors Bright and Cheerful, Luxurious Furnishings, Superior Appointments, Skilled Attendance, First-class Cuisine.



Dr. C. B. BURR, Medical Director,
FLINT, MICH.

RICH RED BLOOD OR BLOOD RICHNESS

Is the main desideratum in many cases. Richness of the circulating fluid in those important basic elements of vitality—hæmoglobin and oxygen.

Pepto-Mangan ("Gude")

INFUSES THIS DESIRABLE RICHNESS IN CASES OF

ANÆMIA, CHLOROSIS, AMENORRHŒA, DYSMENORRHŒA, RICKETS,
BRIGHT'S DISEASE, Etc.,

By furnishing these necessary hæmoglobin-making and oxygen-carrying elements—Iron and Manganese—in a form for almost immediate absorption. Both repeated "blood counts" and clinical experience go to prove this statement.

PEPTO-MANGAN "GUDE" is put up only in bottles holding $\frac{3}{4}$ xi.

Prescribe original packages, Doctor, and thus avoid substitution. NEVER SOLD IN BULK.
Samples and literature upon application.

M. J. BREITENBACH COMPANY, Sole Agents for U. S. and Canada,

LABORATORY,
LEIPZIG, GERMANY.

56 & 58 WARREN ST., NEW YORK.



HYDROZONE

(30 volumes preserved aqueous solution of H_2O_2)

IS THE MOST POWERFUL ANTISEPTIC AND PUS DESTROYER.
HARMLESS STIMULANT TO HEALTHY GRANULATIONS.

GLYCOZONE

(C. P. Glycerine combined with Ozone)

4 IS THE MOST POWERFUL HEALING AGENT KNOWN.

THESE REMEDIES CURÉ ALL DISEASES CAUSED BY GERMS.

Successfully used in the treatment of Diseases of the Nose, Throat and Chest:

DIPHTHERIA, CROUP, SCARLET FEVER, SORE THROAT, CATARRH OF THE NOSE, OZÆNA, HAY FEVER, LAGRIFFE, BRONCHITIS, ASTHMA, LARYNGITIS, PHARYNGITIS, WHOOPING COUGH, Etc.

Send for free 240-page book "Treatment of Diseases caused by Germs," containing reprints of 120 scientific articles by leading contributors to medical literature.

Physicians remitting 50 cents will receive one complimentary sample of each, "Hydrozone" and "Glycozone" by express charges prepaid.

Hydrozone is put up only in extra small, small, medium and large size bottles bearing a red label, white letters, gold and blue border with my signature.

Glycozone is put up only in 4-oz., 8-oz. and 16-oz. bottles bearing a yellow label, white and black letters, red and blue border with my signature.

Marchand's Eye Balsam cures all inflammatory and contagious diseases of the eyes.

PREPARED ONLY BY

Charles Marchand

Chemist and Graduate of the "Ecole Centrale des Arts et Manufactures de Paris" (France)

Charles Marchand,

28 Prince St., New York.

Sold by leading Druggists.

Avoid imitations.

☞ Mention this Publication.

FOR CHILDREN
employ

**"The Safest of all
Modern Antipyretics."**

LACTOPHENIN

PEDIATRICS (March 15, 1897) quotes from a "Report on New Remedies," by Dr. P. Phillip (*Archiv für Kinderheilkunde*):—

"Stein, Strauss, Schlutius, and Riede publish their experience with *lactophenin*. Its sedative qualities, which make it useful, particularly in typhoid fever, are praised by most of its friends. According to Stein it may be used whenever phenacetin is indicated. Because of its relative safety and possibility of longer use it is much superior to phenacetin. Riede calls it a specific in acute articular rheumatism, producing no ear-symptoms or depression of the heart, as does salicylic acid and salophen."

This summary of the therapeutic value of lactophenin has been variously confirmed by clinical reports during the past year from sources widely apart. We note, for instance, in a report on "Treatment of Typhoid Fever in Children," by Dr. F. Gordon Morrill, Visiting Physician at the Children's Hospital of Boston, Mass. (in *Archives of Pediatrics*, March, 1897), that lactophenin is used in that institution, and that 3 to 8 grain doses "are very effective, and do no harm, as far as my experience goes"; lactophenin produces a drop of 3.5° F. in four hours, according to careful records. "Restful sleep may often be obtained in this way."

Descriptive
Pamphlets
Mailed on
Request

C. F. BOEHRINGER & SOEHNE

7 CEDAR STREET, NEW YORK

Lactophenin
is sold by all
Druggists